



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines


Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.


About Google Book Search


Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>




Fr

QE
527
R23







AN
ACCOUNT
OF SOME
GERMAN VOLCANOS, &c.

Speedily will be published,

IN ONE VOLUME OCTAVO,
MINERALOGICAL AND BOTANICAL
TRAVELS THROUGH ITALY;

Written in the German Language by
MR. HERBER:

Translated into English by R. E. RASPE.

Printed for LOCKYER DAVIS.

AN
A C C O U N T
OF SOME
GERMAN VOLCANOS,
AND THEIR PRODUCTIONS.
WITH A NEW HYPOTHESIS
OF THE PRISMATICAL BASALTES;
ESTABLISHED UPON FACTS.

BEING
AN ESSAY OF PHYSICAL GEOGRAPHY
FOR PHILOSOPHERS AND MINERS.
PUBLISHED AS SUPPLEMENTARY TO
SIR WILLIAM HAMILTON'S OBSERVATIONS
ON THE ITALIAN VOLCANOS.

By R. ^{Rich} E. ^{rich} RASPE.

In nova fert animus mutatas dicere formas.

L O N D O N,
Printed for LOCKYER DAVIS, HOLBORN;
Printer to the Royal Society.
MDCCLXXVI.

THE UNIVERSITY OF CHICAGO
LIBRARY
540 EAST 57TH STREET
CHICAGO, ILL. 60637
U.S.A.

05-7-217.M.H.

Rj-st.
804h
7-25-25
11855

[187] REVISED

ADVERTISEMENT.

MANY Philosophers having of late made use of volcanos and earthquakes as undoubted active principles to explain the inequalities of the earth, it is matter of just surprize why the various nature of volcanos, and their productions should be so long neglected. The Author therefore considered the following facts as natural Supplements to Sir William Hamilton's valuable accounts of

b the

vi ADVERTISEMENT.

the Vesuvian eruptions, and hopes they will be received as improvements of useful Science.

There is the less occasion to enlarge upon the advantages of Geographical, Physical, Mineralogical, and Chemical Observations, as it is obvious that, by rectifying the different romantic hypotheses of the earth, they afford Philosophers an opportunity not only to improve the system and classification of Fossils, but what is above any scientific nomenclator, to lay down sure principles for the too-much neglected and expensive art of Miners. Many curious observations to that purpose have
been

ADVERTISEMENT. vii

been made of late in Italy, Hungary, Dalmatia, and Germany, by Mr. Ferber, Baron Born, and Alb. Fortis, in their Mineralogical and Botanical Travels: all which the Author has an intention of publishing in the English language.

b 2

TABLE

1871

July 1st

Dear Mother

I received your kind letter
of the 27th and was glad
to hear from you and
to hear that you were
all well. I am well
at present and hope
these few lines will find
you all the same.

I have not much news
to write at present. I
am still in the same
place. The weather is
very warm here now.
I have not seen any
of the children since
they left. I hope they
are all well.

Yours affectionately

T A B L E
OF THE
C O N T E N T S.

P A R T I.

DESCRIPTION OF THE VALLEY OF CASSELL.

- § 1. *General form, p. 1.*
§ 2. *Sand-stone mountains, which inclose it on one side, p. 2.*
§ 3. *Habichwald, which incloses it on the other side, p. 3.*
§ 4. *Lower hills of this valley described in general, p. 5.*

b 3

§ 5. *The*

* C O N T E N T S.

- § 5. *The calcareous ones are an ancient stratified bottom of the sea, ibid.*
- § 6. *Uncertain on what sort of ground accumulated, p. 6.*
- § 7. *Covered by different modern strata, p. 7.*
- § 8. *Large bones found buried in them at Cassel, p. 8.*
- § 9. *General consideration of these bones, and conclusion that they belonged to an unknown species of quadrupeds, p. 9.*
- § 10. *The calcareous strata covered with old decays of the Habichwald, rolled and washed down to Welhiden, p. 11.*
- § 11. *With a bed of ashes at Weissenstine, p. 13.*
- § 12. *With decays of the Habichwald in many places, ibid.*
- § 13. *Run under different superincumbent mountains, p. 14.*
- § 14. *Conclusion of their high antiquity; inferior to that of the ground on which they are accumulated, but anterior to that of their superincumbent mountains, ibid.*
- § 15. *Level*

C O N T E N T S. 21

- § 15. *Level of the sea which covered or produced them, p. 16.*
- § 16. *The highest hills which inclose the valley, have once appeared above it as so many islands, p. 18.*

xii CONTENTS.

PART II.

DESCRIPTION OF THE HABICHWALD, AND
OTHER MOUNTAINS OF THE SAME NA-
TURE.

- § 1. *They are superincumbent on calcareous ground, and accordingly new mountains,* p. 20.
- § 2. *Raised and accumulated by subterraneous fire,* p. 24.
- § 3. *Their inner structure and beds agree with the nature of volcanic hills,* p. 28.
- § 4. *No rule of specific gravity to be observed in their different strata, which are described,* p. 30.
- § 5. *Their stones and fossils described in general,* p. 32.
- § 6. *The black vitreous rocks found in solid strata superincumbent on calcareous ground,* p. 33.
- § 7. *On*

C O N T E N T S. xiii

- § 7. *On the Krazenberg, p. 34.*
- § 8. *In a calcareous hill near Frankenhäufen, which is an ancient volcanic crater, p. 35.*
- § 9. *These strata, supposed to be lavas, cooled in subterraneous unerupted masses or running erupted ones, p. 39.*
- § 10. *The black vitreous rocks in solid isolated points under the level of the sea, supposed to be remains of the former or nuclei of volcanos, p. 41.*
- § 11. *The black rock in separated straggling fragments very common, p. 43.*
- § 12. *Different constant forms of the black rocks. The prismatical ones called basalties discovered to be lava, p. 44.*
- § 13. *Why have not all lavas a determinate regular and the same form? p. 48.*
- § 14. *Different circumstances under which the different mixture of the lavas is known by facts to be brought into fusion, eruption, and cooling, p. 49.*
- § 15. *Hypothetical explication of the different constant forms of the Hessian lavas, p. 51.*
- § 16. *The*

xiv C O N T E N T S.

- § 16. *The Hessian and many other prismatical basaltes appear in or under the level of the sea, and seem to be lavas cooled by it,* p. 54.
17. *This hypothesis recommended to further examination in Sicily, and some new-raised Islands in the Archipelago,* p. 57.
- § 18. *Description of the Hessian lavas, and their comparison with the Italian ones,* p. 58.
- § 19. *Their different colours,* p. 61.
- § 20. *The black and black-grey coloured are the soundest, appear in different constant forms, and seem to have been but in a state of fusion,* p. 63.
- § 21. *Their substance, which is vitreous, contains different species of sberl,* p. 64.
- § 22. *A new species of crystallized sberl-masses, not to be confounded with granite or granitello,* p. 66.
- § 23. *Queries concerning the origin and nature of the volcanic sberls,* p. 70.

§ 24. *The*

C O N T E N T S. xv

- § 24. *The vitrifications in the hardest lavas described, p. 72.*
- § 25. *The green vitrifications a new species of precious stones, ibid.*
- § 26. *The black vitrifications known by the Ancients, p. 73.*
- § 27. *The spongy lavas never appear in any regular form; and seem to have been erupted running lavas, p. 74.*
- § 28. *They contain in their holes several parasitical aqueous, calcareous, and hornstone congelations, p. 78.*
- § 29. *Of the Vicentine achates enhydros, and the Iceland chalcedonies, produced in volcanic matrices, p. 79.*
- § 30. *The burnt and scorified lavas described in general, p. 82.*
- § 31. *Appear inclosed in tufa, ibid.*
- § 32. *The tarras-stone, a scorified lava, described, p. 83.*
- § 33. *The volcanic ashes of the Habichwald, p. 85.*

xvi. C O N T E N T S.,

- § 34. *A stratum of blueish ashes contains some walker-clay, which confirms Mr. Beaume's theory on the origin of clay,* p. 87.
- § 35. *Several other species of volcanic ashes,* p. 90.
- § 36. *Tufo is produced by ferruminated volcanic ashes,* p. 92.
- § 37. *The τώπος of the ancients, and use of tufo,* p. 94.
- § 38. *The volcanic tufo employed in buildings,* *ibid.*
- § 39. *The tufo described,* p. 96.
- § 40. *Contains different stones,* p. 97.
- § 41. *Class I. Ancient stones,* p. 99.
- § 42. *Class II. Volcanic stones; as ashes, lavas, crystallized sberl-masses, lamellated sberl, sberl-glümmer, and white amiantb,* p. 101.
- § 43. *Class III. Parasitical stones,* p. 105.
- § 44. *Reca-*

C O N T E N T S. xvii

§ 44. *Recapitulation, and conclusion that the Hessian lavas are somewhat different from the Italian ones, p. 106.*

§ 45. *Some general account of other volcanos in Germany, p. 110.*

PART

PART III.

UTILITY OF THESE ENQUIRIES, AND OF
THE VOLCANIC PRODUCTIONS.

- § 1. *General remarks, p. 114.*
- § 2. *Enquiries of this kind improve the knowledge of the physical geography, and the expensive art of the miners, p. 116.*
- § 3. *Use of the volcanic productions, p. 118.*
- § 4. *Use of the basaltes and the harder lavas, p. 119.*
- § 5. *Use of volcanic tarras, neglected in Hesse, p. 123.*
- § 6. *Tarras and volcanic cinders employed by the Ancients in cast vaults and walls, p. 124.*
- § 7. *Example of a cast vaulting of calcareous topbus in Germany, p. 126.*
- § 8. *Volcanic tarras gives an excellent cement, p. 127.*

§ 9. *The*

C O N T E N T S. xix

- § 9. *The puzzolana described according to its volcanic origin, use amongst the Ancients, and its preparation, p. 127.*
- § 10. *The casting of vaults and walls by a puzzolana-cement, used by the Ancients and the Italians, recommended, p. 132.*
- § 11. *The Hessian tarras proved by fact to give an excellent cement under water, and to be equal in goodness to the Dutch tarras, p. 133.*
- § 12. *Mineral waters produced by the volcanic ashes in the garden at Weissenstine, p. 135.*
- § 13. *The use of tufo described before, p. 136.*

PART

THE
[Illegible text]

[Illegible text]

[Illegible text]

[Illegible text]

P A R T I.
D E S C R I P T I O N
O F T H E
V A L L E Y O F C A S S E L L .

§ 1. *General Form.*

THE large valley, wherein the city of *Cassell* is situated, is on every side inclosed by a vast circle of high mountains, and intersected by different lower sloping hillocks.

B

§ 2. *Sand-*

§ 2. *Sand-stone Mountains, which inclose it on one Side.*

The *higher mountains* extend on the South East part of the river Fulde, from Münden in the Electorate of Hanover, Landwernhagen and Spickershausen, to Sichelstein, Wendhausen and Kauffungen; and thence behind Oxhausen and Crumbach to Berghausen, at which place they are called the Soere-Forest, and are cut through by the Fulde.

This whole ridge or chain of mountains consists, as far as I know, of Sandstone-Strata, which are generally horizontal. Never containing any marine bodies, I should be inclined to consider them as *original* or *primogenial mountains*, and to rank them with the pretended *primogenial hornslate-granites* and *metallic mountains*, which in no respect have any better claim to such a denomination, and should rather be called the most ancient mountains, as being generally covered by more modern strata.

Branches

Branches of these sand rocks stretch near Wolfanger and Berghausen over the Fulde, and run, near the former place, Simmerhausen and Knickhagen, to the Reinarts-Forest; but on this side of Berghausen and Freienhagen they are blended with and lost under several lower hills and under the higher Habichwald, as appears by some memorable circumstances described in the sequel.

§ 3. Habichwald, which incloses it on the
Upper State.

The *Habichwald*, whose steep and shaggy summits compose the second or the North-West part of the before-mentioned wide circle of higher mountains, is entirely different from the nature of these sandstones. It is remarkably the highest ridge of mountains thereabout, and, including the Dörnberg, with several mountains of the like nature, it forms a large square of the country, its outline, of above four German or

4 DESCRIPTION OF THE

twenty English miles, running from Weissenstein to Harleshausen, Heckerhausen, Weimar, Zierenberg, Dornberg, Ehlen, Elgershausen, Alten-Ritta, Alten-Bauna, Nordhausen, and Wahlershausen. My readers, who have never happened to be in this country, so much deserving to be seen, and so much signalized by British valour, are referred to an excellent Plan, drawn by Captain Dumont de Venemont, and published with an account of the glorious expeditions of the allied armies after the battle of Wilhelmsthal, which was fought in *June 1762* *. This plan is a model of an excellent topographical map, and will give a distinct idea of the different and relative elevation of the mountains, of their direction, and of the extent of the beautiful valley of Cassel now under consideration.

* Plan de la premiere partie des Operations faites par l'armee alliee contre l'armee Francoise en 1762; to be had at Heydinger's in the Strand.

§ 4. *Lower*

§ 4. *Lower bills of this valley described in general.*

The *lower bills* of the large valley, included by the above-mentioned higher ones, are, on the N. W. side of the Fulde, either of a calcareous nature, or composed of the accumulated decays and rubbish [gefbade] washed down from, and belonging to the surrounding Habichwald,

§ 5. *The calcareous ones are an ancient stratified bottom of the Sea.*

The *calcareous bills*, especially the *Wineberg* and *Krazenberg*, on which the city of Cassel is situated, together with the *Monkberg*, which runs from the city up to *Upper-Vilmst*, consist of parallel strata; dip in different inclinations to the South; run in a N. W. direction, and shew in a certain depth a hard marble-like blueish-grey limestone, which is broken into many vertical fissures, and is filled with some scarce petri-

6 DESCRIPTION OF THE

fications, or rather nuclei of, for the most part, unknown marine bodies, such as cornua ammonis, entrochi, and the like species pelagiæ of plants and animals inhabiting the deepest seas.

Hence it is evident, that these calcareous strata have been anciently the bottom of a sea; but when, or for how many centuries, cannot possibly be guessed. We see however the most evident proofs of its having deposited rubbish and shells. These beds have probably been shaken and split by earthquakes, which have raised and brought them to their present inclinations to the horizon.

§ 6. *Uncertain on what sort of ground accumulated.*

I have not hitherto discovered to full satisfaction on what sort of ground these strata were accumulated, when this part of the continent was covered with sea; but probably it was in some places a sandstone-rock,

VALLEY OF CASSELL.

rock, as one may guess by its before-described neighbourhood, and by several circumstances, which shall be taken notice of in the sequel. In several other places it might be argillaceous slate, which Mr. Ferber* from the Alps to the most Southern parts of Italy has constantly found below the calcareous hills of that country. A stratum of blue clay, discovered and laid open by the brook of Welheiden, in the valley below the Wineberg near Cassel, countenances this opinion.

§ 7. Covered by different modern Strata.

Our calcaceous strata are covered by,

1. A clayish mould (*humus bartensis argillacea*).
2. Yellow ferruginous loam or brick-clay.

* See his Letters on the Natural History of Italy, printed in German at Prague 1773; which will soon appear in the English language.

§. DESCRIPTION OF THE

3. Yellow, greenish, and reddish marle, more or less petrified, which in this country goes under the name of Monkberg, perhaps because the before-mentioned hill of that name has no other exterior mould but this sort of marle. Immediately after these beds, follow the limestone strata.

§ 8. *Large Bones found buried in them at Cassell.*

Foundations of buildings cannot be safely laid in the upper part of Cassell but in orat upon this marle-stratum, which very often puts the builders to great expence; and has occasioned, when that part of the city was first planned out for the French Refugees, a discovery deserving notice. It is said, and there is no reason to doubt it, that they found then in one of the superior clay-strata several bones and teeth, belonging to an elephant, or to a quadruped of a like size, which are kept in the Landgrave's public collection of fossils. The tufts then dug

dug out, according to the dimensions of the several fragments, are judged to be above three feet in length. The grinders are of a proportionable bigness.

§ 9. *General consideration of these bones, and conclusion that they belonged to an unknown species of quadrupeds.*

Bones and teeth of the same kind have been found in the most Northern countries, and also in North America, under the same circumstances; that is, laying in a state of calcination in the uppermost earth-clay and swamp-strata, or in subterraneous caverns, and being never mixed with marine productions, or inclosed in found-stone-beds, there is good reason to believe that such bulky quadrupeds have been in former times, after the sea had retired, indigenous inhabitants of the countries where their remains are dug out. I will not repeat the difficulties, for the most part imaginary ones, which have hitherto attended the explanations

plications of this singular phenomenon; nor will I tire my readers by relating the unnecessary pains taken by the ingenious, to free the question from these difficulties. I would rather refer them to a Dissertation of the celebrated Anatomist Dr. Hunter*, and to another of my own†, concerning a great quantity of such skeletons, which some years ago were discovered near the Ohio in North America. Dr. Hunter demonstrates, that the grinders and several other of these bones do not belong to elephants, but to an unknown species of carnivorous animals; and after some strong inductions against the changes in the inclination of the Ecliptic, which without any necessity, or good evidence, have been supposed to explain this phenomenon, I have hazarded the conjecture, that this species of carnivorous animals, and perhaps a more Northern kind of elephants, has been extirpated, perhaps by the same means as were employed with the

* Philos. Transf. vol. LVIII. † Ibid. vol. LIX.

like

like success against the lions in the Southern parts of Europe, against the wolves and wild boars in England, and against the bears, urus', and elks in Germany, which Cæsar, though he never penetrated further than the Rhine, mentions as inhabitants of Germany*. But I return to our calcareous hills.

§ 10. *The calcareous Strata covered with old decays of the Habichwald, rolled and washed down to Welheiden.*

Beyond Welheiden, on the left of the new-planted green road which leads to Wissenstein, these strata are lost under a gentle sloping ground, which ascends to the Habichwald and the Druselthal, and chiefly consists of the decays and rubbish of the high-towering steeper Habichwald, that thence have been washed and rolled down upon the limestone by the ever-mining waters of a rapid rivulet, called the Drusel.

* Cæsar de Bello Gallico, lib. vi.

[See

[See Pl. I.] This rivulet springs on the very summit of the Habichwald, and, though inconsiderable in dry weather, acquires very often, by heavy rains or melting snows, impetuous force. Under whatever supposition we consider this ground, whether it be accumulated on the limestone strata when still covered with sea, or after the sea had left them bare and dry, its height and large extent obliges us to believe it a work of many ages, and that a long series of centuries was requisite for so small a rivulet to force its way to the depth of a valley like the Druselthal, through the high rocks of the Habichwald, thence to tear and to carry off such an immense quantity of hard stones, to roll these decays on the limestone ground at a distance of three English miles, near to the very gates of Cassel, and finally, to grind and to blunt these stones, which mostly are as hard as flint, into the pebble-like form, in which they appear every where, and especially on the borders of the rivulet, which visibly, even at present,

sent, never ceases in its unwearied course to blunt, by rolling, whatever lyes in its way.

§ 11. *Covered with a bed of ashes at Wissenstein.*

Between Wahlerhausen and Wissenstein the calcareous strata are lost under, and covered with a hill, on which the castle and the *platte forme* of Wissenstein are situated. [See Pl. I.] This hill is a mass of a spongy, fine and grey bluish mould like ashes, washed down from the higher and immediately connected Habichwald. It cannot be ranked among the clays, marles, and sands, since it has not their requisite toughness, cohesion, effervescence, and pal-
pability.

§ 12. *Covered with decays of the Habichwald in many places.*

Near Kirchditmold the calcareous strata ascend gently up to the Habichwald, and disappear under its ruins.

On

12. DESCRIPTION OF THE

On the side of this high forest they baffle out behind Upper-Vilmar, near Willemsthal, Calden, Franckenhäusen, Burg-Uffelen, and Grabenstein.

§ 13. *Run under different superincumbent mountains.*

Behind the Habichwald and Harleshausen they appear again from under its decays, ascend to the isolate Dornberg, whose foundation is entirely calcareous, and run by the way of Zierenberg and West-Uffelen, in the country of Waldeck, beyond the Diemel, in the Bishoprick of Paderborn, and thence into the Forest of Teutoberg, in the county of Lippe.

§ 14. *Conclusion of their high Antiquity; inferior to that of the grounds on which they are accumulated, but anterior to that of their superincumbent mountains.*

I have several times travelled over these parts, and after repeated examination found,
that

VALLEY OF CASSELL. 15

that their limestone strata, though now and then interrupted and covered by large valleys, being throughout of the same colour and substance, and containing the same sorts of petrifications or nuclei of conchæ, and species pelagiæ, are of the same kind. This gives me a right to conclude;

1. That all these parts of the continent have been covered with a sea for as many ages as might be requisite to produce, at its deepest bottom, so great a variety of calcareous strata.

2. That these strata are of a newer date than the above-mentioned sandstone, or argillaceous slate strata, which lie deeper, and are [See Pl. I.] of the highest antiquity.

3. That they are anterior to the decays of the Habichwald [See Pl. I.]; and,

4. Even of a more ancient origin than the Habichwald, the Dornberg, and several other superincumbent hills of that kind. [See Pl. I.]

I am

I am not inclined to lose myself in a nearer examination of their antiquity, which my readers are desired to fix as they please, according to the wants or to the advantages of that system of Chronology, which they have a mind to, or are convinced of.

§ 15. *Level of the Sea, which covered or produced them.*

There is more hope of success and satisfaction in determining the level of the sea, which once covered or produced these calcareous strata, or this ancient sea bottom. It is traced on the sides of the higher surrounding hills of the large valley of Cassell, by a stratum of sea mud, which in several places I have examined myself, and according to repeated mensurations found to lie in all these places on the same elevation. [Plate I.]

This remarkable stratum is to be met with in the Sneckenberg within the garden
at

at Wissenstein [Plate I.]; and on both sides along the descent of the Habichwald; near that place. I saw it likewise on the other side of the river Fulde, at a distance of seven or eight English miles, in the descent from Wendhausen to Kauffungen; and it occurs again at a still greater distance, that of at least ten English miles from Cassell; near Deuten, Bessé, and Gudensberg. In all these places it is at the same horizontal elevation as marked in the Section Plate I. and of the same nature. It consists of a friable, and more or less indurated ferruginous marle or mud, of an ochraceous yellow-brownish colour; and it contains an innumerable quantity of calcined littoral-shells, such as *Chamas crassas majores læves æquales polyginglymas*; *Conchas cordatas læves majores & minores*; *Pectines auritos minores*; *Turbines*; *Cochleas*; *Strombos minores*; *Entalia*; *Tubulos vermiculares*; and finally, some small Coral-fungites — that are known species and common on many shores

18 DESCRIPTION OF THE

shores of the Northern seas and strong evidences that an ancient sea covered that part of the Continent at this level. It was perhaps the same sea, which produced the above-mentioned calcareous strata and species *marinas pelagias*; and having left so many evident littoral-traces on the sides of the Habichwald, one might seem entitled by that circumstance to suppose, that the Habichwald, and the like sort of mountains, which are accumulated on a marine limestone ground, have been produced in the midst of that ancient sea, long before it retreated.

§ 16. *The highest hills, which inclose the valley, have once appeared above it as so many Islands.*

The highest summits of the higher mountains, which inclose the valley of Cassel, must accordingly needs have been above the level of this ancient sea as so many islands;

VALLÉY OF CASSELL. 19

islands; but especially the Habichwald, the Dornberg, and several other hills on the N. W. of the river Fulde, which, from their structure, and the nature of their fossils, I conclude to be a sort of mountains entirely different from all the before-mentioned supposed original arenaceous and secondary calcareous ones.

PART II.

DESCRIPTION

OF THE

HABICHWALD, AND OTHER MOUNTAINS OF
THE SAME NATURE.

§ 1. *They are superincumbent on calcareous ground, and accordingly new mountains.*

I Come to a more minute description of the Habichwald, and other mountains of the same nature. They are superincumbent mountains (Aufgesetzte Berge) surrounded on all sides by lower calcareous hills, on which they are accumulated. This, though

though obvious in many other places, is more clearly to be seen and to be evidenced by the Dornberg, which to the middle of its elevation is entirely calcareous, and leaves no room for any conjecture of such hills being original or primogenial mountains.

They cannot be considered neither as accumulated by the action of the sea (Flotz-Geburge), since they are destitute of the parallel-strata of marine mountains: and though now and then there appear in them something like parallel-strata, there is never found inclosed in their substance any thing like adventitious organic marine productions.

Nor can they have been accumulated or deposited by rivers and brooks, since there is no higher ground, whence they could have been supposed to flow; and since the slime or mud of rivers and brooks are nowhere found to produce rock so hard and vitrescent as theirs.

It would be equally hazardous to attempt explaining their origin by pretended inundations, covering and overflowing their highest summits. I shall not insist on the deficiency of water; and I will grant for a moment, that the fountains of the deep, and the windows of heaven, or an auxiliar comet's tail, and other such miraculous unvidenced and improbable hypotheses, could have once drowned the world. But would that give them their vitreous nature? Inundations can be supposed to produce, by the depositions of their sediments, a variety of parallel strata, which does not appear in the mountains we treat of; but they never can have composed wild vitreous rocks, in that situation, form, and quality, in which they appear in the solid and large masses, or huge pointed tops of black rocks, belonging to the highest elevation of the Habichwald, Dornberg, and others.

But perhaps they may have been accumulated on the limestone in these unweildy,
uncouth,

unicouth, enormous masses, by successive sediments of hot wells, or incrustating waters; or produced in the same manner as the ground near Radicofani, a description of which I have laid before the Royal Society at London*; or deposited as the calcareous tophus, or osteocolla-stratum, near Ziierenberg in Hesse, at the very foot of Dornberg, is daily increased; or raised in the same manner as the white saline marble and alabaster hills seem to have been composed †. This conjecture may be supported by the calcareous strata on which they are superincumbent, and which possibly here, as commonly in the places before-mentioned, might produce such incrustating wells and waters; but it is absolutely inconsistent with the vitreous nature of the mountains and masses under consideration, since all sediments, precipitations, and incrustations, of hot wells and waters, are of a calcareous or selenitic nature, effervescent

* Phil. Trans. Vol. LXI. † Ibid. l. c.

by the action of the acids, and easily brought to calcination by common fire.

§ 2. *Raised and accumulated by subterraneous fire.*

These considerations and circumstances do not leave open any hypothesis for explaining their origin, but the conjecture: "that these singular superincumbent mountains are the immediate work of subterraneous fire."

It is fact, that subterraneous fire, and its many successive eruptions, have raised or heaped together the still-burning Mountains *Ætna* and *Vesuvius* on the limestone-strata in Sicily and Italy*, accumulating both these mountains to an elevation, and to an extent, which even surpasses that of the *Hambichwald*. It still continues to work in the volcanoes in Iceland; and there is no good reason to deny the possibility of other Eu-

* Ferber's Letters.

repeat volcanos, situated between Iceland
 and Ætna, and burning in former times.
 There have been found of late many extinct
 volcanos in Italy, stamp'd with visible marks
 of ancient burning, though never spoken of
 in history. Why should not Germany then,
 as well as Italy, afford phenomena of the
 same nature? The sea, which covered these
 parts, as many others in the continent, will
 not, I hope, be alledged against this suppo-
 sition, and thrown upon it to quench the
 German volcanos; since the still-burning
 volcanos are generally situated in the midst
 of the sea, in islands, or near the sea coast,
 and even seem to want sea-water to raise,
 and to support their very flames. It
 would be unfair to conclude, or to cavil any
 thing against their former existence, from
 the silence of history; because ten thousand
 things may really happen every day, and
 pass unnotic'd; and German history, in
 respect to the long series of former forgot-
 ten ages, begins but from yesterday—from
 Cæsar,

Cæsar, Drusus, and Germanicus, our generous conquerors; or from S. Bonifacius and Charles the Great, our ungenerous apostles. In this light I certainly am allowed to venture that hypothesis, which not only is possible, but even seems to be a necessary supposition, since, besides the above-mentioned insufficient natural causes of superincumbent mountains, there have been to this time no others known at all.

This supposition, like many others, might be possible, and seem necessary; and nevertheless be improbable, may prove entirely false. But it has rather every quality of historical demonstration, as standing upon firm unshaken foundations, and good authorities, which are;

1. The inner structure of these mountains; and,
2. The nature and quality of their stones and fossils.

Both

Both agree with the nature of the Italian volcanos, which, though observed and described by many curious and ingenious men of every age, have not been sufficiently described by any mineralogist. But by Mr. Ferber.

This ingenious gentleman was, in the year 1768, when I saw him first at Cassel, neither sufficiently acquainted with, nor a friend to, the volcanic system; accordingly he did not then entirely agree with my opinion. But since the year 1771, when he visited Italy, he has made my opinion his own, and confirmed it by his valuable mineralogical letters on Italy, which are as good evidence for me as the Hessian volcanos, whose nature I shall now describe, and which lie open, together with Mr. Ferber's Letters, to every one, who may want still fuller instruction, and have a mind to a nearer examination.

§ 3. *Their inner structure and beds agree with the nature of volcanic hills.*

The inner structure of the Habichwald, of the Dornberg, and of many hills of that kind in Lower-Hesse, may be easily observed, wherever they are steep and shaggy, or wherever snow-rain- and brook-waters have laid them bare, in valleys, natural cascades, and ravins. There is the same facility in several quarries, but especially in the subterraneous galleries and aqueducts, which, on account of the famous cascades at Wilfenstein, have been dug in the Habichwald, near and under the octogon of the Carlberg.

The Habichwald, being by far the greatest and highest mass of this kind of mountains, and having been less degraded and changed by rain, wind, and weather, deserves a peculiar notice in respect to its inner structure. It shews a variety of accumulated

cumulated vegetable mould, ashes, clays, marles, sands, black vitreous rocks, flags, and other stones, under different inclinations to the horizon, and in strata different from those which water produces, or is supposed to have produced, being never so exactly separated from among themselves, but commonly indiscriminately and insensibly blended together, crossing one another in their direction, and dipping on every side, in almost every sense, often in parallel inclination, to the sloping and descent of the mountain, often falling exactly against the mountain and its general direction. This singular position, nature, and variety of stones and strata, cannot be easily explained by the common orological hypotheses: but it ceases to be a problematical riddle as soon as we suppose this vast mountain to have been a burning volcano in former times, and to have accordingly, at different times, and by different craters, vomited sand, ashes, brimstones, flags, and lavas.

30 DESCRIPTION OF THE

§ 4. *No rule of specific gravity to be observed in their different strata, which are described.*

As the strata of this mountain have no regular direction, dipping, and form, they are destitute likewise of any rule of specific gravity. At the entrance of the Druselthal, and at the foot of the Habichwald, to the right of the road from Cassell to the coal mines, there is a large stratum of light spongy ferruminated albes, below a stratum of very ponderous and hard vitreous black stone. On the N. E. side of Weissenstein there do appear, on the descent of the Habichwald, in several places, and in different elevations, immense masses of tufa in different large beds, covered at the summit of the mountain by heavier black stones. The castle of Weissenstein is situated ;

1. On a foundation of spongy friable mould or albes, buried behind the great basin, which is higher up the hill,

2. Under

2. Under a very heavy and compact mass of basalt. [See Plate I.] Higher upwards,

3. By clay and sand, in isolated nests or lumps [Plate I.]; and,

4. By great globular masses of black vitreous stone. Immediately under the octagon, at the top of the cascades, offers,

5. A very large and extensive solid stratum of black stones, whose broken fragments are scattered and straggling all along the descent of the hill, as marked in Pl. I. Higher, and on both sides of the octagon, follows,

6. An extensive stratum of light spongy slugs, which to the left of the octagon dips in the mountain, under

7. A stratum of heavy black rock, that runs all along the long-stretched summit to its steep descent in the Druselthal, and the valley of Cassel.

The

32 DESCRIPTION OF THE

The same irregularity in respect to the specific gravity of the strata of ashes, tufo, and lavas, is observed in Monte Somma, Vesuvius *, and other still burning volcanos; and it is similar to their irregular vomiting of ashes, sands, and fluid lavas.

§ 5. *Their stones and fossils described in general.*

The stones and earths of the Habichwald, and its kindred mountains, or the substance of their strata, which have been described, are,

1. Black vitreous rocks of different weight, solidity, and form.
2. Tarras, and flags.
3. Tufo.
4. Different species of stones, inclosed in the black rocks, and the tufo.
5. Ashes.

* Ferber's Letters.

6. Clay

- 6a. Clay and marle.
- 7. Sand.
- 8. Vegetable mould. And,
- 9. Finally, a vast bed of coals, lying in a valley, gently sloping from the very summit of the Habichwald, but according to many circumstances, independent from its rocks, to which it is superior, and of a more modern and different origin; for which reasons I forbear to speak of it in particular, intending only to describe the nature and respective situation of the before-mentioned fossils, which are related to each other.

§ 6. *The black vitreous rocks, found in solid strata, superincumbent on calcareous ground.*

The black rocks are called in Hesse *Swarze-Wacken*, and would go in the mineralogical systems under the general name of horn-stone.

They are found either, 1. in solid, found, and coherent strata; or, 2. in solid ridges
D and

and isolated points; or, 3. broken and scattered in their ruins.

If found in solid strata, they are now and then found immediately superincumbent on the solid calcareous strata, which I have described before as lying at the bottom of the valley of Cassell, and of the Habichwald. They ought to be very obvious in that situation on the Dornberg, which up to its middle is calcareous, and at its summit is all over covered with fragments and masses of this rock. I have not, however, remarked on this mountain any place, where this situation might be clearly observed. Therefore I shall describe two other places of that kind, which I have had the good luck to find elsewhere.

§ 7. *On the Krazenberg.*

The first is on the Krazenberg, in a limestone quarry, near an intrenchment raised in the last war over against Kirk-Ditmold. The calcareous beds are there running from
the

the S. E. and the city of Cassell to the opposite Habichwald; and in a line, which is parallel to this higher mountain, there goes across the run of the limestone a large vertical vein or fissure (similar to what the German miners would call a *gang*), which is filled with black rock. It is spungy and full of holes; but these holes are, for the most part, filled with white calcareous spar, occasioned by the neighbouring limestone.

§ 8. *In a calcareous bill near Frankenhäusen, which is an ancient volcanic crater.*

The second place here to be mentioned is still more remarkable, and to be found on a small limestone-hill near Frankenhäusen, bordering on the road from Cassell to Carls-häven. I owe its discovery to the penetrating look of his Excellency the Minister and Lieutenant General Baron de Slieffen. On the top of this hill is a conical crater, whose circumference is about three hundred feet. It is broken through the almost

horizontal strata of marle and limestone; and what is extremely singular, it has been on its inner sides all around incrustated and hanged with a covering of grey and black spungy rock, which, where it is broken, appears to have been one foot in thickness. [See Plate II. n. 1, 2.] This vitreous covering, incrustation, or tapestry, hangs flat upon the borders of the crater, and before the broken reddish and whitish marle and limestone strata. A cast stucco-floor, sunk down in a conical crater before its full petrification, would produce something like this phenomenon, because the black stony incrustation sticks every where close to the sides of the crater, has yielded to their inequalities, which it expresses by its bending bunches and concavities, and is cracked and broken in many irregular but coherent pieces. The same sort of vitreous black and spungy rock is found at the bottom of this crater in large irregularly-split masses, and every circumstance cooperates to bring

to mind what Mr. Ferber * says, in his description of the Italian volcanos: "that very often their funnels are covered but by a thin covering or floor of indurated lava, which often gives way and falls down to their bottoms, probably filled with nothing but masses of lava." And certainly we might not be wrong in looking upon this crater as a funnel of a small volcano; 1. on account of the vitreous nature of the black rock, which covers its sides, and fills its bottom, and will, in the sequel, be proved to be a lava; 2. because the strata of reddish and whitish marl and limestone, to which this vitreous covering sticks, are visibly affected and somewhat calcined, by the heat of this lava; 3. because a running lava issued from this crater, as may even yet be seen by its beginning within the crater, [Plate II. n. 3.] though the outside of the hill, and the current, I speak of, be entirely buried at present under vegetable mould, and does not appear, like many other

* Letters, p. 149. 188.

volcanos, to be externally clad and covered by ashes or volcanic tuff—a circumstance, which may have been occasioned by having been for a long time exposed to the washings of decomposing rain or inundations, or even by having been produced under the level of the ancient sea, which I have traced before, and certainly was higher than its summit.

This remarkable and instructive natural curiosity is perhaps the only one yet known of that kind. However it is likely to be demolished in few years, since they have begun already to remove a good deal of the inner covering of lava, for paving the Carlshaven-road, which goes near to it; and if the Landgrave do not stop this demolition, as may be hoped, it will be soon lost for ever. At all events, and to give it its due celebrity, I have it drawn on Plate II.

§ 9. *These strata, supposed to be lavas, cooled in subterraneous unerupted masses, or running erupted ones.*

The large extensive strata of the black rock under consideration are, for the most part, in a nearly horizontal situation, and thus to be seen at different elevations of the Habichwald. They are besides, without any respect to specific gravity, in different places, now under, now above other lighter strata of ashes, tufo, and slags. I beg leave to consider these solid rock-strata as lavas, cooled in large extensive melted masses under ground, or cooled after their eruption, as large running lavas. I shall prove it in the sequel. The greatest stratum of this kind, or rather the most extensive lava, which I had an opportunity to observe in these parts, is at the top of the Carlsberg in the Habichwald, under the foundations of the octogon. [Plate I.] Behind the grotto of Polyphemus and on its both sides,

D 4

especially

40 DESCRIPTION OF THE

especially to the right, it appears bare and open on the sloping of the steep mountain. It is there about fifty feet thick. I could not distinguish from what crater it came, or where it directed its course, if it be at all to be considered as an erupted running lava. But its length or extent is at least 270 German fathom, or 2170 Rhinish feet, this being the length of a subterraneous gallery, which, on account of an aqueduct and the cascades, has been cut with great expence, and a true Roman perseverance, through this hard vitreous solid rocky mass. Its length and extent may be by far more considerable; and perhaps all the solid vitreous black rocks on both sides of the octogon, and on the whole summit of the Carlberg, together with the romantic heaps of its dreadful ruins, rolled down its steep descent into the lower valleys, might be justly considered as belonging to the same lava. It is covered near the cascades with flags or tarras-stones, and these by a smaller lava or bed of black rock, which to the left of the octogon

octagon dips under ground in a direction towards the old Winterkasten.

Thus far of the black rocks in solid strata,

§ 10. *The black vitreous rocks in solid isolated points under the level of the sea considered, as remains of the former, or as nuclei of ancient volcanos.*

Their solid ridges and isolated points, standing free and bare on all sides, are very common in Lower-Hesse. Such is the Katzenstein and the Hohlstein under the Dornberg, between Hapschausen and the village called Dornberg; a peak in the Wildenberg near Wolfhagen; the castle-hill at Felsberg; the castle-hill at Aldenburg; the Maderstine near Gudensberg, and another of that kind near the road thence to Cassell. Most part, if not all of these bare ridges and isolated points, are below the before-mentioned level of the old sea, and there is generally something very particular in the form

form or breaking of their rocks, which shall be noticed afterwards. Having ventured to call the solid strata of the black rocks cooled lavas, in erupted runnings, or unerupted melted subterraneous masses, a denomination which their situation and many other qualities will justify; I venture, in the same hope, to consider these ridges and isolated points as remains of old running lavas, or as kernels and nuclei of old volcanos, deprived by rain, snow, overflowing water, and a long series of destroying centuries, of their external covering of ashes, cinders, and sands; which seems the more probable, as in their neighbourhood, and even at their feet, for example at the Holefine, their own ruins of black rock, and great masses of tufa and indurated ashes, are to be met with.

§ 11. *The black rocks very common in separated stragglings fragments.*

The black rocks, broken and separated from their natural place, are very common in several steep descents of the Habichwald, in the Druselthal, to the right hand of the octagon near Harleshausen, on that part of the Carlberg which lies between the cascades and the Moriz-grotto, [Plate I.] behind the Paul's hermitage, on the Dornberg, and in a hundred other places, where their cinereous or tuff ground giving way and being washed of great masses of the superior rock-strata tumbled down in wild ruin; or where the before-described bare ridges and isolated points mouldering away, were dissolved to ruins, rolled and washed down into the vallies. They are in every situation, in solid strata and bare ridges, as well as in their ruins, so very common in those parts, which I have had opportunity to see, that generally all the roads and cities of Lower-Hesse have no other pavement.

§ 12. *Dif-*

§ 12. *Different constant forms of the black rocks. The prismatic ones, called basaltes, discovered to be lavas.*

The *form and figure*, which they spontaneously affect when broken, is very different. I have on this account observed the following varieties.

1. Their form is *indeterminate* in the great lava-stratum on the Carlberg, [Plate I.] as clearly appears in the subterraneous gallery, and in many fragments, dispersed every where in the valleys and ground immediately under its steep summit.

2. They appear in *irregular slates or tables*, and in a *vertical position*, near the old Winterkasten on the Carlberg.

3. In *large globular masses*, composed by concentric and less compact coats, and containing in the middle a remarkably hard and compact globular nucleus or nucleus, they are obvious in a large stratum near the
Moriz.

Moriz-grotto behind the Sneckenberg, and in different other places at the Carlberg*.

4. They are *irregular polygons*, and tending to a prismatic columnar form, in a stratum behind the great basin up to the Moriz-grotto at Wiffentine, at the level with the Sneckenberg. [Plate I.]

5. In a *columnar prismatic regular form* they appear in large masses or strata of close-lying columns in the castle-hill at Felsberg and Aldenburg, in the Maderstine near Gudensberg, in the Widelsberg near Wolfhagen, in the Holestine under the Dornberg, and, as I have been told, but never observed myself, in the Dornberg.

These prisms or columns are of a quadrangular, pentagonal, hexagonal, and heptagonal form; flat at the top; and in the same stratum, or rather the same mass of

* Similar masses have been found by John Strange, Esq; F. R. S. in the Euganean and Vicentine volcanic hills; their description and drawings given in his account, Phil. Trans. vol. LXV.

the same length : but, the length of these different masses being different, they differ likewise in length from six feet to twenty, and above. They have a thickness from ten to fifteen inches. To give an idea of their wondrous form and masses, I refer my readers to the engravings of the castle-hill of Felsberg and of the Maderstine, which are annexed to their description in the Philosophical Transactions for the year 1771. Till very lately, the origin and nature of these singular stones was considered as inexplicable ; but a great many correspondent observations, and a nearer examination of their substance and situation, has, since the year 1768, convinced Mr. Desmarest*, myself †, and Mr. Ferber ‡ ;

a. That the substance of these figured prismatical and columnar vitreous black

* Explication des Planches de l'Encyclopedie. Livraison cinquieme, 1768.

† Phil. Transf. vol. LXI. German Dissertations of the R. Soc. at Goettingen, tom. I. 1771.

‡ Letters, 1773.

stones,

stones, is to be considered almost every where* as a ferruginous mass, impregnated with much iron, and smelted by heat or fire.

b. That, appearing in prismatical forms, it is named, by the Ancients and the Mineralogists, basaltes.

c. That, in an indeterminate form, it goes in Italy under the name of lava, or felce and pietre dure.

d. That by refrigeration it gets a prismatical form, as most part of smelted ores and metals get by cooling a determinate crystallized form in their substance †.

* Ferber's Letters, p. 64. 272, 273.

† One of the most singular evidences of this common phenomenon happened some years ago at Clausthal in the Harz forest. The smelting of a high- or roast-furnace, alternately filled with coals, fluor, and pulverised arsenical lead- and silver-ore, being interrupted by accident, the ore cooled and appeared sticking in it in large coagulated loose and spongy blue-glittering masses, all over covered on the surface with quadrangular crystallisations, in a form which the French call à la Grecque, unknown whether ever any where else produced by art or nature.

Fuller

Fuller evidences of this hypothesis shall be given in the sequel.

§ 13. *Why all lavas have not a determinate regular and the same form?*

But why have not all our black rocks and all the Italian lavas the same and a constant determinate prismatical form? Even the prismatical columnar lavas or basaltics show varieties of forms. The Irish basaltics in the Giants-causeway in the county of Antrim, appears in polygone atticulate columns or prisms. There ought to be some natural reason for that, whatever it be: Is it owing to the difference of their substance and mixture? to the different sloping of the ground on which these fiery melted masses run forward? to the different quickness of their motion? to their different fluidity? or is it owing rather to the manner of their refrigeration? All these circumstances may, without any doubt, have influenced their different forms, but our black rocks are prismatical.

prismatical basaltés being of the same mixture and substance, seem to indicate that their different fluidity, fusion, motion, and refrigeration, have been the proximate causes of their before-described determinate and indeterminate forms and figures.

§ 14. *Different circumstances under which the different mixture of lavas is known by facts, to be brought into fusion, eruption, and cooling.*

Whatever be the cause of the fusion and fluidity of the smelted lavas, whether it be a strong heating fermentation, as seems highly probable, or whether it be a subterraneous fire, as commonly is supposed without sufficient evidence, it is a fact, wanting no evidence, that the lavas of *Ætna* and *Vesuvius* either break their way from under ground through the lower sides of these mountains, or ascend like boiling water to the very brim of their craters, which they overflow at last, and thence run

E

down

down into the lower vallies; or that they are vomited and thrown out of these craters in fluid burning lumps. It is fact likewise, that the lavas appearing in their eruptions like fiery torrents, either overflow the continent above the sea, and refrigerate there; or that, no eruption ensuing, they refrigerate again within the volcanos; or finally, that sometimes, according to the situation of the ground, which they overflow, they run down into the sea or other water. Nor can there be any doubt, that by similar causes violent fermentations, fusions, nay eruptions of smelted substances may be produced even at, and under, the very bottom and level of the sea. Whoever reflects on the great number of islands and high volcanic hills, with boiling and smoking of the ocean produced in the deepest seas, will be less amazed at the boldness and singularity of this supposition.

§ 15. *Hypothetical explication of the different constant forms of the Hessian lavas.*

Now in all these different cases, the cooling and refrigeration of the lavas must needs be very different.

Undoubtedly they cool slowest in the holes and craters of the volcanos, where their mass is less troubled by motion, and their heat less influenced by air and water.

They are sooner refrigerated in the torrents, running after their eruptions over dry land, as being influenced by more cooling circumstances of wind, weather, rain, or snow; sooner still, if thrown out in fluid or burning lumps; but soonest and most violently in those fiery torrents, which run into neighbouring seas or other waters, or even break their way immediately into the sea.

Though I am neither inclined, nor able and willing to explain the mode by which

these circumstances give a different form to the hot-fluid or burning masses of lava, they certainly had some share in it, as every one will agree with me, that different degrees of heat or cold, by means entirely unknown to us, visibly influence the crystallisations of salts. Accordingly the external circumstances, under which the different forms of lavas are found, seem to entitle me to the following conjectures, relating to the different modes of refrigeration, and their influence on the masses of lava.

The *undeterminate forms* of the black rock or lava-stratum at the summit of the Carlberg, seem to be such, because the lava is at that place in an almost horizontal situation, which kept the whole mass undisturbed, and gave it an opportunity to refrigerate very slowly in itself.

The *irregular broken slates* lying with the former above the level of the sea, which I have traced before, [Plate I.] at the very summit of the Habichwald, seem to belong

to a lava cooled by air, and perhaps by rain or snow, which split it, as heated glass cracks if suddenly brought in contact with cold water.

The concentric globular lumps, with a hard solid nucleus in their middle, are never found in a solid stratum, but here and there scattered about and heaped together. They seem to have been thrown out as fluid hot lava-masses, and falling through the air, and rolling on the descent of the mountain, to have acquired a globular form. The supposition that their external crust suddenly cooled, by the ambient air, while they moved fast, and that they were rolled afterwards over beds of ashes and small volcanic stones, which commonly cover the sides of the volcano, accounts very well for their hard nodules, which, unaffected by the air, cooled slower. It accounts likewise for their external cracking and breaking into concentric scales, which are of a less compact substance, resembling

rather ashes ferruminated with small volcanic stones. The origin of the phialæ Bononienſes and cracking glaſſes is in ſome reſpect like to this hypotheſis, and Mr. Ferber * gives nearly the ſame explication of the ſame phenomenon in Italy.

§ 16. *The Heſſian and many other prismatic baſaltes appear in or under the level of the ſea, and ſeem to be lavas cooled by or under it.*

The regular prismatic columns black rocks or baſaltes ſeem, according to their ſituation and quality in this country, and in many others, to be lavas, which in a hot ſtate ſtand broke their way from under ground immediately in the ſea, or run into it, or cooled under its level, without any eruption, within the ſtrata, caverns, and holes, wherein they had been brought to fuſion.

All our Heſſian baſaltes, which I have examined, are viſibly under the ancient level

* Letters, p. 278, 279.

of the sea, which I have traced before. [Plate I.] This remarkable circumstance is evidenced to conviction, by a large bed of lava, which is throughout inclining to a prismatical form, and is situated immediately under the Moriz-grotto at Wissenfins, exactly at the level with the Sneckenberg, which contains a stratum of calcined littoral shells. [Plate I.] The castle-hills at Felsberg and Aldenburg, together with the Maderfins near Gudensberg, are undoubtedly below the same level.

2. Above this level I never have found, neither on the Habichwald nor on the Dornberg, any prismatical basaltic mass. These facts agree with the following in other countries.

3. The Irish basaltic in the Antrim Giants-causeway, which are articulate prismatical columns, together with those of the same kind, which Mr. Banks has of late discovered in the island Staffa *, are situated

* Pennant's Voyage to the Hebrides,

near or even in the marine level, may far stretching into and under the sea. Those at Dunbar in Scotland * are in a similar position; and the gigantic basalt-pillars near Lauganes in Iceland, which are nine feet diameter, are continually washed, and for a great part overflowed, by the sea †. Ronca, a hill in the Veronese territory near the Valle del Buso, consists of alternate marine and volcanic productions, among which a bed of black lava split in prismatic forms clearly appears to be covered by a calcareous stratum filled with a variety of marine petrifications ‡, which greatly countenance, in my opinion, Bolca, a hill, likewise in the Veronese, is of the same mixed marine and volcanic kind, consisting in its summit of calcareous slate, famous for its fine petrifications of marine fishes, and covering an immense variety of lavas and other volcanic productions §. The prismatic basaltes near

* Pennant's Tour to Scotland.

† Drawings in Mr. Banks's collection.

‡ Ferber, p. 53. § Ibid. l. c.

Bolzena,

Bolsena, between Monte Fiascone and Aquaspendente in the Roman State, are nearly in the level of the lake of Bolsena*.

4. The most exact observers of Vesuvius; Sir William Hamilton at Naples; and Mr. Ferber, have never met with any prismatic lava or basaltic in its many old and new erupted lavas, which cooled in the continent, and above the surface of the sea; nor do I remember to have found them mentioned in Sir William Hamilton's, or other descriptions of the gigantic Ætnean lavas, which like large rivers have overflowed the land.

§ 17. *This hypothesis recommended to further examination in Sicily, and some new volcanic Islands in the Archipelago.*

However, I recommend this singular phenomenon, and my bold hypothesis, to the

* Ferber, l. c. p. 381.

examination and judgement of Messrs. Desmarest, Fortis, Born, Ferber, Sir William Hamilton, and Mr. John Strange, as the most competent judges, hoping that if the cooling under ground and the sea level, or in sea or other water, has really had any share in the prismatical form of basaltic or lavas, more evidences of it ought to be found in many volcanic islands, but especially in the too much neglected new raised islands of the Archipelago* and in the large Sicilian lavas, which near Catania and many other places reached the sea.

§. 18. *Description of the Plesian Lavas, and their comparison with the Italian ones.*

Whatever I have hitherto said concerning the situation and the forms of our black vitreous rocks, or of refrigerated lavas, is but hypothetical. Therefore the question naturally arises: whether this may be confirmed by their nature and by their com-

* In the year 1763. described and recommended to philosophical observations in my *Specimen hist. naturalis de novis insulis. Amstelodam.*

parison

parison with true lavas? and whether our black vitreous rocks, basaltes, and the known lavas, be one and the same sort of stones, and accordingly of the same origin? Every one of my readers has full right to put this question to me, who, unsatisfied by hypothetical solutions, chooses rather to *see* than to *hear*. Whoever has no opportunity to convince himself by ocular inspection, may be convinced by Sir William Hamilton*, and by Mri Ferber †: "that lavas and *most part of the prismatic basaltes do not differ in themselves, but only in their forms.*" and I desire these gentlemen to believe me on my word, that our prismatic basaltes at Felsberg and elsewhere are, on account of their colour, substance, and mixture, of the very same mass as our hardest black and grey rocks, which

* See his letter of Dec. 22, 1769, which he did me the favour to write from Naples, inserted in the German Dissertations of the R. Soc. at Goettingen, vol. I. p. 92.

† Letters, p. 65, 66. and 279.

have

have all the qualities of many Italian and other lavas, and which, together with the basaltcs, for that reason, I not only henceforth shall constantly call lavas, but presently describe, with convenient exactness, an account of their various inner qualities. I would not however be charged with the imputation as if I did, without exception or proper allowance to facts, declare every sort of black vitreous rocks, together with every species of their kindred hornstone-rocks and prismatical basaltcs, to be produced by subterraneous extinct fire or fermentation. I am convinced by facts and led by good authority to presume, that Nature works similar effects by different causes. It is but lately that I have learned by Mr. Ferber, though in the main he be of my opinion, that there are some scarce species of basaltcs, which seem to have been produced by water *, and that there are at Wilkissen in Bohemia, as in different other places,

* Letters, p. 272.

rhomboidal

rhomboidal clay-slate and granite columns, which owe their crystallisation to water*. I intend here only to speak of our black rocks, which in themselves and through all their varieties show incontestable marks of fire and fusion, are undoubtedly nearly related to our basaltes, and resemble the Italian and other lavas.

§ 19. *Their different colours.*

Our lavas are like the Italian ones, which Mr. Feber has so scientifically described in his Letters†, of a different colour, solidity, weight, texture, and appearance, according to their different mixture, to their degree of fusion and combustion, and to their refrigeration.

They are of a black, black-grey, ash-grey, and reddish brown and ferruginous colour.

* Mineralogical Account of Bohemia, German edit. p. 122, 125.

† P. 175.

62 DESCRIPTION OF THE

The two former forts are most frequent. The basaltic at Felsberg, Aldenburg, and Maderstine, are black or black-grey.

The ash-grey coloured are in different shades to be found in the high borders of a deep rivulet near a village called Bodiger, and in dispersed straggling fragments between Upper-Vilmar and Burg-Uffelen.

The reddish and brown-coloured are very scarce, and but in straggling pieces and fragments to be found at the foot of the Holestine under the Dornberg. This and the ash-grey species I never have happened to find completely solid, found porous, and in a prismatical form, whence it seems to follow, that they have been in a state of conflagration, which supposes an eruption, and has greatly lessened their weight and solidity.

§ 20. *The black and black-grey coloured are the soundest and hardest; appear in different constant forms; and seem to have been but in a state of fusion.*

The black and black-grey lavas, which are the soundest, hardest, and heaviest, are probably such lavas as have been just in a state of perfect fusion, and cooled within themselves, or on a sudden. For this reason and on account of their prismatic-columnar, concentric-globular, slate-like or undeterminate forms, they seem to be different from those which, having been in a state of eruption and conflagration, never offer any such regular form or sound solidity. They are generally susceptible of a fine polish, and withstand for many centuries the action of wind and weather, and even exterior violence in the public roads. Though extremely hard and compact, they are however but slowly dissolved or affected by air, since exposed to it for many centuries

64 DESCRIPTION OF THE

turies they get a white-greyish colour, which now and then is found penetrating for a tenth part of an inch into their harder and deeper-coloured inner substance.

§ 21. *Their substance, which is vitreous, contains different species of sberl.*

Their substance is a fine mass, without any visible grain; vitreous, but without any remarkable gloss on the fracture. It is scarce if ever entirely pure, homogeneous, and of the same mixture of colour, being more or less mixed and internally blended.

1. With deeper-coloured spots of the same substance;
2. With small whitish, green, and blackish, often scarce visible glass or sberl sparks or flakes;
3. With larger masses of mixed white, yellow, green, and blackish crystalized sberl,

sherl, commonly appearing in the form of blunted larger pebbles;

4. With black lamellated sherl; and now and then, but very seldom,

5. With yellow, deep grass-green, and black vitrifications, harder than flint.

The fonder and harder black and grey lavas of Vesuvius, described by Mr. Ferber *, are generally of the same mixture as these under consideration, except that they contain a greater quantity of sherls in larger crystals, and in a greater variety of determinate forms †, and that they never contain any of the above-described mixed crystallized sherl-masses, N° 3.

* Letters, p. 164. 167.

† Such as a polygone, globular, garnet-like, and hexagonal pointed or truncated columnar, and other forms.

§ 22. *A new species of crystallized spherulites, not to be confounded with granite or granitello.*

Here it may be questioned: "whether these pebble-like mixed crystallized spherulites of our harder lavas, may be at all ranked amongst the spherulites? or, whether they be a particular undescribed species?"

When I first attempted an examination of our basaltes and lavas, I was very uncertain what to make of these masses. According to external appearance, which in mineralogical objects is hardly ever to be depended upon, I considered them as a species of granite fragments, which I supposed to be thrown out with the lava; but as they are really destitute of the quartz, feldspath, and mica-particles, the essential

* See my account of the Hessian basaltes, in the German Dissertations of the R. Soc. at Goettingen, tom. I.

ingredients

Ingredients of granites *, I readily gave up this erroneous opinion, and declared them in the year 1768, when I was acquainted with Mr. Ferber, and on his authority, to be a tin-ore, similar to a species of zinn-

* I know very well that some Italians call granitello the speckled and coarse substance of the basaltes in Monte Rosso, which is exactly resembling that of the large basalt-pillars near Langanefs in Ireland, as may be seen in Mr. Banks's collection, and even has some likeness with the granite-like Hessian spher-masses; *sed verba valent sicut nummi*; and I could wish to see the pretended Euganean columnar or other granitello chemically and closely examined by so excellent an observer as Mr. John Strange, in order to give into the hypothesis (Phil. Trans. vol. LXV. part. I.); and consider this granitello as a granite, and the Monte Rosso columnar basaltes as a granite-bed melted and crystallized locally, or in *statu quo*. Monte di Diavolo near S. Giovanniignano is, according to Mr. Ferber's account (Letters, p. 66), as many other basaltine and lava-masses, superincumbent on calcareous strata; and granites have never been found inferior to them, being besides substantially different from the volcanic spher-concretions which I was speaking of, and which seem to be common in the Monte Rosso basaltes.

wald in Bohemia*. However we were equally wrong in this conjecture, which I have been convinced of by several chemical assays, by a closer examination of their matrix and its concomitant circumstances, but especially by Mr. Ferber's late descriptions of the Italian spherl and lava species.

These mixed crystallized spherl-masses agree with the known spherls in their melting very easily by the blow-pipe, and in their giving to the borax-glass a deep green colour. They do not contain any tin. Though congealed in smaller crystals than the common Italian lava-spherls, they are found accompanied by unequivocal lamellated spherls, inclosed in the same mass of lava or basalt. Like the Italian larger spherl-crystals they dissolve in the air, and moulder in an ochraceous, ferruginous, argillaceous powder. Accordingly, they are to be considered; " as a particular un-

* L. c. and described by Mr. Ferber in his mineralogical account of Bohemia, p. 132.

“ described

“ described species of spherls, peculiar to
 “ our Hessian lavas or basaltcs *.” If these
 spherl-concretions be found inclosed in the
 sounder and harder lavas or basaltcs, they
 are tolerably compact and remarkably pen-
 derous; take a polish inferior to that of
 their lava- or basalt-matrix; an indifferent
 polish to be had on their yellow and faint-
 green crystals, but a splendid one on the
 black and dark-green ones, which are re-
 markably hard, and strike fire with steel.
 Sometimes they are found in tufa, ferru-
 minated in ashes with different stones and
 lava-pieces, which appear to have been
 thrown out by volcanic eruptions, and then
 they are observed to be sometimes wrapt up
 in a mass of spongy, slaggy, and burnt
 lava; which shows them to have been
 thrown out with this burning lava from the

* It is sprinkled, but in smaller patches and lumps in
 the Langanefs basaltcs in Iceland, in Mr. Banks's col-
 lection, and it seems to be common in the basaltine
 masses of Monte Rosso, as described by Mr. John
 Strange.

deeper volcanic funnels, and to have been either in a state of fusion, or covered with burning lava lumps.

§ 23. *Queries concerning the origin and nature of the volcanic sherls.*

They never have offered themselves to me in our tarras or those lavas that have been scorified, though they very often contain a brittle and glossy lamellated sherl. This remarkable circumstance naturally occasions the following questions :

1. Whether the substance of these mixed white, yellow, green, and black sherl-masses, be the fixed phlogiston of our lavas and basaltes?
2. Whether this phlogiston be congealed into crystalline lumps by the suffocation or cooling of the fluid lavas? or,
3. Whether these masses, whose substance is by fusion thoroughly mixed with the mass

mass of the Italian and our lavas, be found in the deepest volcanic rocks and funnels as peculiar pre-existent rocks, or be produced and inclosed like pebbles in the cooled lavas of former times?

4. Whether they be thrown up, melted in the fluid lava, and by the intensest degree of heat, which consumes even lavas to slags, they be volatilised either wholly or in part, or changed and vitrified to glass-frittas and brittle and lamellated sherls, such as are found not only in the sounder, but likewise in the scorified, lavas and basalts?

There is no circumstance to warrant the supposition of their having been like other parasitical stones generated and crystallized by water in holes and bladders of solid lava; but future chemical assays will certainly throw light on these queries, and demonstrate the origin of lavas and of these and other sherl-species, which Mr. Ferber is inclined to ascribe to subterraneous fire.

§ 24. *The vitrifications in the hardest lavas described.*

The *vitrifications* or *frittas* of our lavas seem to be of the same substance as the *flint-masses*, since they stick to the same volcanic matrix, and have the colour of their dark-green and black crystals, and of the lamellated *flint*; but they are remarkably harder, strike fire with steel, cut glass, and might with justice be ranked among the precious stones. They are of a deep grass-green, a yellow, and a black colour.

§ 25. *The green vitrifications a new species of precious stones.*

The *green ones* are of a deep grass-green hue, have a pebble-like blunt form, and are as it were cast in the mass of lava or basalt. Their hardness is remarkable; resists the English file, and requires in grinding and polishing the same art as the hardest

est precious stones, according to the trials and evidence of the court-jeweller Hefs at Cassell. But it is pity that they are so scarce, and commonly cracked and split. Else their fine colour and hardness would entitle them to be ornaments of the rich, as well as other ornamental coloured natural glasses, which on no better account have been ennobled by prejudice and superstition; and if they should be found destitute of their fabulous magic craft, they certainly would not appear to be as resplendent and as electrical.

§ 26. *The black vitrifications known by the Ancients.*

The *black vitrifications* or *frittas* are to be found in as undeterminate forms. However, I have a specimen in a pyramidal form, with blunted angles externally rifled like prismatical or stangen-spherl. Their inner texture is now and then similar to that of the lamellated spherl, but they are much harder.

harder. The Ancients knew these vitrifications under the name of *petra obsidiana*. They have a fine black colour; and on account of their hardness, which strikes fire and resists the file, they take a fine polish.

Though I have likewise found in our solid lavas and basaltes *yellow vitrifications* like topaz, they never occurred to me in a size sufficient to assay their hardness, polish, and other qualities.

All these vitrifications offer themselves likewise in the Vicentine and Vesuvian lavas*, though on account of form their species are different from ours.

§ 27. The *spongy lavas* never appear in any regular form; and seem to have been erupted from running lavas.

So much of our solid lavas and basaltes, and their admixed corall stones. I proceed to a second species, to the *spongy*,

* Forber, p. 173.

Skummy, and honey-comb lavas, whose nature gives visible evidence of their fiery origin. They never have offered to me in a prismatical-slate or concentric nucleus form. I have found them immediately laid on the limestone-strata, and commonly in straggling fragments; but never in sound strata. Where they appeared stratified and coherent, they might be supposed belonging to erupted lava-torrents. Mr. Ferber*, speaking of the slaggy Vesuvian lava, says: "that during the eruptions, it runs out of Vesuvius with violent ebullition and skimming, which by means of the inclosed air produces its various-sized holes and bladders, and makes its superficies very porous." On this authority our spongy honey-comb lavas are to be ascribed to neighbouring volcanic craters, or to the surface of lava-torrents; and in fact, they are found in the above-mentioned calcareous crater near Frankenhäusen; [Pl. II.] and on both ends of the gallery, or the

* Letters, p. 173.

large

large lava-bed on the Carlberg. [Pl. I.] Their mass, mixture, the glass and shreds in them, are the very same as those in the solid lavas, from which they differ in nothing but, 1. the want of a prismatical or other determinate form; 2. their lesser weight; and, 3. their air-holes. I have found the following varieties.

1. A black ponderous species with round holes, in straggling pieces in the Druselthal.

2. Of the same colour, straggling under the Holesine on the Dornberg, with lengthened irregular holes.

3. A black species in the inner part scorified and burnt, but externally solid though slaggy. Frequent in tufo; to be more minutely described afterwards.

4. A grey species, with large air-holes, dispersed on the public road between Upper-Vilmar and Frankenhausen.

5. Another

5. Another grey species, to be found in the calcareous crater near Frankenhausen. Its air-holes small.

6. Another grey-coloured, with smaller holes, in the limestone-quarry on the Krazenberg, described before.

7. A black sort from the lava-bed and the gallery under the octagon on the Carlberg.

8. A whitish-grey, sandy, coarse species, from the borders of a rivulet near Bediger, with large round bladder-holes.

9. A reddish friable; and,

10. A red brownish solid but scarce species, with large holes, under the Holestine on the Dornberg.

§ 28. *They contain in their holes several parasitical aqueous, calcareous, and hornstone congelations.*

The air-holes or bladders of all these lavas are seldom found empty, being commonly filled with stones and earth, which have been deposited and congealed therein by steams or water, and for this reason are of a more modern date than their lava-matrix. You meet often with such holes, which have been but partially filled and incrustated; and some there are, in which these congelations have dissolved and have almost disappeared. The red and brown species, 9. 10. have their air-holes slightly incrustated with ochraceous and brown iron-dust. The whitish-grey from *Bediger*, n. 8. contains some empty bladders, some coloured with brown iron-dust, some filled with glass kopf or button-ore. The holes of the black and grey ones are generally filled with white, and sometimes crystal-
lized

lized calcareous spar; but in the species n. 1. and 2. from the Druselthal and the Holfelthine, they are incrustated with a white milk-coloured, opaque, hard and thin crust, which seems to be flinty and vitrescent, since it resists the action of acids.

§ 29. *Of the Vicentine achates enhydros, and the Iceland chalcidones, produced in volcanic matrices.*

Perhaps Nature intended here to produce chalcidony by the same way as it generates this fine stone in the volcanic ashes and sands near Vicenza*, or in similar volcanic matrixes

* Ferber, p. 21. They are found in nodules, which sometimes are hollow within, and filled still with the water that had produced them. A very remarkable phenomenon! They are known in Italy under the name of *achates enhydros*, and the grain of chalcidony and agathe is certainly not to be distinguished. Abbé Fortis at Venice gave me the following account of them in a letter of January 18, 1772. "L'Achate enhydros, que je vous ai envoyé, se trouve dans la pierre
" brune

80 DESCRIPTION OF THE

matrixes in Iceland, whence lately have been brought stratified large chalcedony-masses, equal in goodness, but far superior in bigness, to those of the East-Indies*. I have in my collection of fossils specimens of these chalcedonies, which convince me

“ brune-volcanique dont je vous ai aussi fait passer un
“ échantillon. On la trouve dans une colline près de
“ Vicence, qui donne aussi sur le sommet & tout au-
“ près de cette pierre même des corps marins en pierre
“ calcaire. Vous ne me demanderez plus à quel prix
“ cette curiosité, est chez nous lorsque je vous aurai dit,
“ que c'est moi même qui les trouve & les fais travail-
“ ler. On a deux ou trois pour cent globules d'Achate,
“ qui ayent de l'eau; & on ne trouve pas quatre pour
“ cent entre ceux qui en ont, dont on peut faire
“ usage.”

* As the court-jeweller Mr. Kefs wanted large pieces of chalcedony to finish the prospect of Rhinisch in Florentine-Musäico or pietre commesse, which by order of the court has been undertaken many years ago, and will be when finished one of the most remarkable pieces of workmanship of that kind. I procured them from Copenhagen by Mr. Danz at Blankenburg near Saalfeld, one of our best merchants for natural curiosities.

that

that their matrix is a sort of volcanic tufa, and that accordingly chalcedony is henceforth to be considered as a parasitical stone or concretion in volcanic ashes or lavas. Who ever thought of that in former times! and who for that very reason can help wishing that henceforth the too-much neglected lavas and volcanic productions, and especially their parasitical concretions, might be subjected to a closer examination?

The lavas hitherto described seem, on account of their external appearance and circumstances, to have been just in a state of fusion; and their weight and solidity give some hopes that a proper chemical analysis will determine their constituent parts, and show out of what species of metals, stones, salts, and earths, the great chemical fire of Nature did prepare and melt them. Mr. Sauffure at Geneva, and Mr. Erxleben at Goettingen, have proposed to undertake these assays, and undoubtedly their pains will not be unrewarded.

§ 30. *The burnt and scorified lavas described in general.*

The *burnt and scorified lavas* have undergone different degrees of combustion. Though coloured like those before-described, which have been but just in a state of fusion, they have lost, by longer and intenser heat, and perhaps by their own conflagration, their solidity, hardness, and weight. They are changed either to stuggy light stones, or to different sorts of ashes. The former, I mean the scorified stones, are found either mixed with ashes and other stones ferruminated into volcanic tuff, or in a large stratum near the octagon on the Carlsberg, which consists of spungy light tarras-stone.

§ 31. *Appear inclosed in tuff.*

The *burnt lavas contained in tuff* have been less consumed by fire, being black or
grey

grey lumps, whose harder and spungy outside shows still the common sherl-spots, masses, and vitrifications; but in the inner parts they are so exactly resembling iron-scoria, that nobody can possibly doubt their former fluidity and burning. They have been perhaps thrown out in burning lumps, extinguished by the ashes in which they fell, but internally consumed by continued intense heat and glowing.

§ 32. *The tarras-stone, a scorified lava, described.*

The *tarras-stones* are as cinders throughout scorified, burnt, spungy; and accordingly very light. They resemble the small reddish cinders in the Roman-Puzzolana; though these commonly be less spungy; but they are entirely similar to the *tarras-stone* from Bockenlime and Andernac, which are very well described and chemically analyzed in the Magazine of Hanover, Nov. 21, 1774. They are found in a large stratum

near the octogon superincumbent on the large solid lava-bed. The surface of the running lavas is very similar to its nature. P. La Torre, giving an account of the Vesuvian eruption in 1731, says: "As the lava advanced, a great many stones tum-
 " bled from its surface. It was covered all
 " over with stones, which were either in
 " their natural state, or calcined and re-
 " sembling burnt tiles or iron-slugs. Their
 " gravity was very different." The former fluidity of our tarras-stones or slags appears on their surface, and in their form, which is extremely various, and sometimes resembling to rifled coral-branches. Mr. Ferber found, on the inner sides of the crater of Vesuvius, a lava which had the same form *. Their surface is covered and ferruminated sometimes with small volcanic stones; but nothing proves their ancient fluidity more clearly than their spongy substance and air-holes. Their colour is ex-

* Letters, p. 176.

ternally

ternally grey, red-brown, or ochraceous, and seems to be occasioned by water soaking through the superior beds of mould or lavas, incrustating them with dissolved ashes, clay, and red ochraceous stalactites. If broken, the fracture is grey or black, like the colour of the solid lavas, from whence they sprung. They never contain the crystallized sherl-masses of the solid less consumed lavas; but sometimes they offer a resplendent brittle lamellated sherl in black sparks and a thick-lamellated spar, which does not effervesce with acids, and appears but very seldom.

§ 33. *The volcanic ashes of the Habichwald.*

The *ashes*, being the remains of the consumed lavas, tarras-stones, or scoriz, are thrown out by the volcanos either pure, or mixed with different sorts of sands and stones, affected or unaffected by fire, fluid, or burning; which, according to their different mixture, causes different beds of

ashes, that are somewhat resembling to the aqueous or submarine strata, as appears on Vesuvius, and in the tuffo-quarries on the Habichwald.

These ashes, being first loose, must needs roll and slide down from the summit of the volcanos, if they be steep, or they might be washed down by rain-water into the adjacent vallies. By such accidents the cities of Herculaneum, Pompeia, and Stabize, have been entirely buried; and for similar reasons we find our volcanic ashes on the foot of the Carlsberg still loose and pure in the flat ground, on which the Sneckenberg and the castle of Wissenstine are built. They could not remain on the volcanic summits in their first beds, but where they found horizontal grounds. Accordingly they appear in large horizontal strata in the tuffo-quarries on the Habichwald behind Paul's-hermitage, in the Druselthal, and at the summit of the hill above a small village called *Hof*.

I have

I have observed the following varieties of ashes.

§ 34. *A stratum of blueish ashes contains some walker-clay, which confirms Mr. Beaume's theory on the origin of clay.*

A blue-grey and friable species stands in a horizontal bed under the palace of Wifsenline and the Sneckenberg, as [Pl. I.] can be observed every where, especially in the terraces under the palace and under the bowling-green in the gardens. These ashes are pure, without any mixture of other volcanic stones, perhaps because they had been situated under the level of the old sea, [Pl. I.] and washed by it or been brought down from the mountain by rain- and snow-water. This species does not effervesce with acids, and is friable to an impalpable powder, for which reason it cannot be ranked amongst the finer sands. In open air or in water it dissolves and melts without any coherence, for which reason it cannot pass for clay. It consists

of fine grey and black points, and is mixed with some fine talc or mica flakes; it appears by the microscope. Its elixivation properly evaporated gives some salt, which for its effervescence seems to be alkaline. The ochraceous sediments of the waters springing from or soaking through this stratum indicate some vitriol and iron, and its sulphureous smell the presence of a sulphureous acid, which probably occasions in this stratum some phenomena strongly confirming Mr. Baume's theory and experiments on the origin of clay, as produced by vitreous earth saturated by sulphureous acid. The phenomena I am speaking of, are great lumps of a fine tough walker-clay (*Wallerii*; Spec. xxix.) and straggling shale-nodules, both now and then involved in this cinereous stratum without any mark of their being adventitious to the place.

The *walker-clay* is of the colour of the ashes, but somewhat ferruginous. It moulders in open air into lamellated slates,

and requires an efflorescence of the same salt, which by elutriation can be extracted from the ashes. Its particles are fat and saponaceous to the touch, getting by rubbing a polish on the fractures. They effervesce with acids, which may be owing to the mixture of the alkaline salt.

The *marle-nodules* resemble the *ludi helmontii*, and split into scaly pieces, commonly joined together by a crystallized yellow spar: but the least exterior violence separates them again, and then the incrustations of these scaly pieces appear like fine yellow velvet, spotted here and there with pyrites. Their effervescence with acids is not only an evidence of their having been produced by the before-described walkercay, but likewise of an admixed alkaline substance.

§ 35. *Several other species of volcanic ashes.*

Whitish-grey ashes friable and extremely light, are found above the tuff-beds behind the Paul's-hermitage at Wissenfene. They are mixed with fine lava-sand and sherl-points, of a white, yellow, ferruginous, and black colour. They resist acids,

A *blue species* equally friable and mixed with black sherl-points, is found in the same place.

Grey and brown-coloured ashes mixed with a great number of small volcanic stones, cinders, and lamellated sherl, resisting acids, are found in a large bed in the Druselthal underneath the Habichwald.

Future enquiries may perhaps discover more sorts of ashes in these volcanic countries, nay there is some probability that several earth- and clay-strata in the vallies about

about the Habichwald and Dornberg are owing to ashes in some places changed by sulphureous acids. The coal-stratum on the summit of the Habichwald may be perhaps ascribed to the same cause as being in a great many respects different in its nature and situation from other coal-beds. But not having yet fully examined these objects, I cannot support this conjecture, except by Mr. Ferber's striking confirmation of Mr. Beaume's theory on the origin of clay, and by his remarkable account* of the solfatara near Naples, and the alum-works at Tolfa in the Roman dominions. The ever-working sulphureous acid has in these places not only changed volcanic ashes and lavas into clay, but even imparted to it an aluminous nature †.

§ 36. *Tufo*

* Letters, p. 187. 243.

† The same may possibly have happened in the famous coal-pits on the Wiffener, in the remarkable clay-pits and in the rich alum-works near Allmerode in Hesse. I have observed in these parts a great many
stones

§ 36. *Tufa is produced by ferruginated volcanic ashes.*

Volcanic ashes, if pure and ferruginated, have produced the above-described cingereous friable stones; but thrown out or washed together with various stones, affected or unaffected by fire, fluid and infused, they have produced the large and various tufa beds, which according to Mr. Ferber's Letters (p. 144, 145, 146.) are frequent in the neighbourhood of volcanos, because they hardly ever cease to throw out

stones resembling those of the volcanic Habichwald, such as lime and sand-stones, and harder lava-pieces; and in the same tract of land, that is to say, between the Fulda and Weser, I have examined eighteen years ago a coal-pit and alüm-work belonging to Baron Hanstein, and situated on the Steinberg near Münden, where he also was excited by effluvia from the slag and the ashes of brown coal or fossil wood, which was found there unpetrified and greatly impregnated with pyrites and sulphur. The coals of the Wiskener are of the same kind.

ashes

ashes and stones. The Paupilippo and other Mills near Naples are only large heaps of such tuff, which ought to be carefully distinguished from the different calcareous tuffus or osteocolla precipitated by water. Strata of this cinereous tuff are very common round about the Vesuvius. Herculaneum, Pompeja, and Stabizæ, are buried under them * ; and the Habichwald in its whole and vast circumference is but an accumulated heap of alternate tuff and lava, as appears by the frequent large and horizontal tuff-beds between the Druselthal and the palace of Wissenstine, behind Paul's hermitage in the gardens belonging to it, but especially by the quarries on the summit of the mountain near Hof, and in many other places. The Dornberg was of old likewise clad with tuff, as appears by its large masses and lumps scattered at its feet near the Holesstine, and in different other places of its vast extent.

* Ferber, p. 145. 184. +86.

§ 37. *The τώρος of the ancients, and use of tufo.*

The ancients employed tufo in the walls and vaults of their buildings *, but if the late Abbé Winckelmann by the authority of Pausanias (lib. v. c. 10.) intended to prove, that the temple of Jupiter at Elis was constructed with tufo, he certainly was wrong, because the τώρος, which Pausanias speaks of, was, according to Theophrastus de Lapidibus, on account of its colour and solidity, like the Parian marble, which bears no resemblance at all to the volcanic tufo.

§ 38. *The volcanic tufo employed in buildings.*

There is no doubt that the ancient inhabitants of Italy employed this sort of stone in the inner mass of their walls and in their

* Remarks on the architecture and buildings of the ancients, p. 2—4.

larger buildings *. Even the present inhabitants have not discontinued that practice, which is equally in use in some parts of Germany, because this stone is very common, very easy to be cut into convenient forms, to be found in large pieces, and never cracked and split. The famous cascades, grottos, and the octogon at Wissenstine, stupendous works, which would bear a comparison with the gigantic vast Egyptian buildings, and the greatest pieces of architecture of antiquity, are built with this stone. They would have lasted through many future ages, eloquent monuments of the magnificence and taste of the late Landgrave Charles of Hesse-Cassel, who built them, if their architect Guarneri had lined or covered them on the outside with a more solid and durable stone, since all the volcanic tufos, and especially the Hessian ones, moulder away if exposed to air, rain, and frost. This seems to be owing to the want of a sufficient

* Letters, p. 278, 279.

quantity

quantity of ashes, and even to a change of these ashes, deprived perhaps of their cementing and ferruminating quality by sulphureous acids and damps, when thrown out with so many burning stones. There are some circumstances to support such a conjecture. Exposed for some time to rain and water, and even in the quarries, they appear spongy, and their holes are filled or incrustated with whitish clay or marle, the result of ashes and sulphureous acids. Exposed to fire, even to the stronger kitchen or furnace fire, they are lasting without decay, a quality of argillaceous stones.

§ 29. The tufa described.

The colour of our tufa is different, as its mixture and solidity; but commonly all its varieties are found in different beds in every quarry opened on the Habichwald. There is,

1. A blackish-grey;

2. A

2. A bluish-grey; and,

3. A yellow species; and every colour of different mixture and solidity. If they be of a fine grain, the inclosed lavas cinders and sherls, being very small, appear in the mass of the stone like different strata, and make the stone resemble the product of water. The yellow species resists the air less than any other.

§ 40. *Contains different stones.*

In the grosser-grained species are to be observed, mixed and ferruminated with ashes, a great variety of stones of different bigness, from the smallest point to a very considerable size; and these deserve a closer examination, as giving evidence and conjectures about the nature and quality of the volcanos which threw them out, and about their deepest rock-strata, to which some of them did belong in former times. Therefore I shall describe all their different spe-

H cics,

cies, as observed by me in the walls of the cascades and elsewhere, and rank them in different classes.

The *first class* is of such stones as are still raw and unaffected by fire, justly supposed to be thrown out and to be separated from the deeper stone-beds, produced by water and belonging to pre-existing and perhaps original or primitive mountains.

The *second class* is to contain the lavas and the products of fire, that is, such stones as have been pre-existing in other forms, but have been by fire and heat affected, melted, changed, or consumed.

The *third class*, which Mr. Ferber has not noticed as all in his volcanic accounts, shall contain the newer parasitical stones which have been in the tufo or lavas produced by water, soaking through their cracks holes and strata.

§ 41. Class I. *Ancient bones.*

1. *White sand-stone lumps*, pure and snow-white; now and then spotted with small black iron-points; the grain extremely fine; their hardness, consistence, and solidity, different; some friable to a fine sand, which on account of its white colour seems to give a good sort of glass.

2. The same sort of *sand-stone* externally vitrified into a hard yellow vitreous crust, which strikes fire with steel. It is scarce in the tufo, and then but in smaller pieces; but common in large straggling blocks and smaller among the ruins of the Habichwald near the palace and the ponds of Wissenstein. [Plate I.] Their nature, situation, and appearance, prove, 1. that they have been affected by fire, and even brought to vitrification; 2. that a species of white straggling rocks, called in these countries *white wacke*, is but a more or less vitrified sand-stone; 3. that the strata under the

Habichwald of a fine sand-stone are perhaps a continuation of the sand-stone hills on the other side of the Fulde, which are called the Soere, and have been described before. Its existence under the Habichwald is still more evidenced by a ridge of large sand-stone rocks near the palace of Wissenstine, which appear above the cinereous ground on which it is situated. Their white shining colour is said to have occasioned the name of this place; and their romantic form, embellished by convenient plantations of shrubs and trees, is one of the most picturesque objects of the noble garden which belongs to this palace. The fragments of white sand-stone, dug out with black solid lava, of an old subterraneous gallery under the old or the smaller Winterkassen behind Paul's-hermitage, and the sand-pits near the Moriz-grotto behind the Sneckenberg, together with,

2: *Reddish sand-stone*; and,

4. A like-coloured species with mica, both very common in tufo, had probably no other origin.

5. *White marle lumps*, effervescent with acids, are likewise inclosed in tufo, but scarce. They belong probably to the calcareous ground which surrounds this mountain, and on which it has been accumulated.

6. *Grey burnt clay*, resisting the acids, very common in tufo.

7. *Red*, and

8. *Yellow ochraceous burnt clay* is found in smaller fragments, which commonly break into concentric pieces. I have not hitherto observed in this tufo true calcined limestones, which in respect to the inferior calcareous strata should seem to be very common, and which Mr. Ferber has found scattered about Vesuvius, and inclosed in the Vesuvian tufo- and lava-beds.

§ 42. Class II. Of volcanic stones, or immediate productions of fire, inclosed in the ash- and tuff-stones of the Habichwald.

9. All the above-described species of ashes, being the ferruminating cement of tuff;

10. Fragments of solid lava of different size;

11. Ditto spongy and honey-combed;

12. Ditto half burnt and extinguished, in the inner parts consumed and burnt to cinders;

13. Crystallized sberl-masser as described before. They are very often wrapped in a spongy or grey lava, fluid when thrown out.

14. Lamellated sberl;

15. Sberl-glimmer; both in black lumps, remarkably large and ponderous.

16. A species of *white amiantb*; has longitudinal fibres; smooth and saponaceous to the touch; scarce coherent; loose; giving way to the pressure of the fingers; light; for the most part swimming on water; resisting acids, and resembling by its fibrous texture and lightness burnt wood entirely consumed to ashes. It is found in the tufa quarries in large but commonly flat pieces, wrapt in a ferruginous ochraceous covering.

Though possibly our ashes and tufos might, as well as the Italian ones, contain burnt wood*, and other adventitious organic bodies †, and therefore this substance be considered as the cinereous remains of burnt wood; there are several circumstances which forbid such a conjecture. Its snow-white colour certainly does not answer the idea of burnt wood, which in the Italian or

* P. La Torre, Hist. of Vesuvius.

† Ferber, pt 126. 184—186. 237.

Herculaneum ashes and tufos* rather resemble charcoals, and its tender coherence and substance is as inconsistent with such an opinion. But all these qualities, together with its fibrous texture, do exactly agree with the amianthine fossil-cork (*Suber montanum Wallerii*, *Spec.* 148.), which is observed among the volcanic stones of Vesuvius †, and according to Mr. Needham's observations nearly related to the lightest cinders, consisting only in an iron calx, entirely deprived of its phlogiston ‡. An intense volcanic fire might probably occasion such an operation. Whether our *amianthine-cork* might be by furnace-fire brought to a black vitrification like that species which Wallerius has described, or whether it might like other species of amianth and asbest, resist the furnace, and

* La Torre, l. c. Winkelmann's Letters on the Herculaneum discoveries, 1762, p. 13. 63.

† Ferber, p. 159.

‡ Mem. de l'acad. R. des Sciences, 1760.

yield

yield only to the burning mirror, is to be tried by future chemical experiments.

§ 43. Class III. *Or the parasitical stones produced by water or steams in the substance of our tufo, contains,*

18. *Whitish and yellowish clay deposited in its spongy holes;*

19. *White calcareous spar, which has incrustated and filled these holes, very common in the tufo-quarry behind Paul's-hermitage in the gardens at Wiffenstine; of a fine grain; very compact, but less than the horn-flinty depositions of the Vicentine schates or rather chalcedonius enhydros; less likewise than the similar horn-flint concretions in our sounder lavas, which I have mentioned before.*

20. *Calcareous spar, of a yellow and whitish colour in large crystals, found in straggling tufo-pieces near the crater at Frankenhäusen;*

21. *Ochra-*

21. *Ochraceous iron-ore*; breaks concentric; light; resembling some richer button-ores; found in lumps, like indurated clay in the tufo-quarry behind Paul's-hermitage,

§ 49. Recapitulation and conclusion that the Hessian lavas are somewhat different from the Italian ones.

Thus far of the stones and earth-species, which have occurred to me in the solid beds of the Habichwald, the Dornberg, and other ridges or isolated conical hills of that kind. These I have been obliged to consider as volcanic, 1. since, being evidently accumulated and superincumbent on marine beds, they cannot be ranked among the pretended original or primitive mountains; 2. since their structure, beds, and fossils, do not agree with any known species of modern mountains but the volcanic; 3. since they agree in most parts of their

their qualities with the nature of many Italian volcanos and volcanic productions; and since, 4. they manifestly show marks of subterraneous fermentation, heat, eruption, and fire, in their structure, as well as the vitreous slaggy nature of their fossils. I have seen, and observed, without prejudice. This I am conscious of, and will I hope appear to every one, who is willing and able to see without prevention. According to this my impartiality I have freely acknowledged, "that in the lavas of Vesuvius and its neighbourhood, no prismatic columnar lavas or basaltics have been observed; and that our lavas differ from those which Mr. Ferber has examined near the Vesuvius and other unequivocal volcanos in Italy, on account of their peculiar and hitherto unobserved crystallized surt-masses." And with the same impartiality I acknowledge still further, "that the still burning volcanos in Italy, Vesuvius, and Solfatara, offer several phenomena, which are not to be

5

" found

108 DESCRIPTION OF THE

“found or observed in our extinct volca-
“nos.” Such are, besides several fossils
described by Mr. Ferber *,

1. The hot wells and steam-baths in the
neighbourhood of Vesuvius †;

2. The suffocating damps and steams
(moffette) of several volcanic tuff-caverns
near Vesuvius †; and,

3. The yellow native sulphur;

4. The red arsenic;

5. The green vitriol;

6. The salmiac; and,

7. The alum; which in the inner crater
of Vesuvius and Solfatara are sublimated
and deposited §. But all these differences

* Letters, p. 158, 159—184.

† Ibid. p. 187. 199—203. La Torre Hist. 69 Ves-
suvius.

‡ La Torre, l. c.

§ Ferber, p. 181—184. 192—196.

intitle nobody to conclude, " that our
 " Hessian basaltes, black rocks, tufe- and
 " tarras-stones, are neither lavas nor of a
 " volcanic origin." The only conclusions,
 which fairly may be drawn from these cir-
 cumstances, and in which I perfectly agree,
 are, " that our volcanic stones, lavas, ba-
 " saltes, and tufe-stones, are, on account,
 " of their substance and mixture, some-
 " what different from the Italian ones * ;
 " and that our volcanos, extinguished and
 " entirely refrigerated so many centuries
 " ago, cannot possibly produce the steam-
 " sulphur and salt-sublimations, which are
 " entirely depending on the action of still-
 " burning and operating subterraneous heat
 " and fire in Vesuvius and Solfatara."
 Their containing the requisite materials for
 many, if not for all these sublimations, ap-
 pears by their description, and will appear
 still more by their future chemical analysis.

* The Iceland lavas and volcanic production (and
 who will doubt their origin?) are certainly as much if
 not more different from the Italian ones.

III DESCRIPTION OF THE

§ 45. *Some general account of other German volcanos.*

Thus we may be without any predilection to the volcanic system convinced, that the Habichwald and the adjacent Dornberg are volcanic mountains. Their remarkable elevation, their large extent of at least 20 English miles square, and their present exterior appearance, seem clearly to indicate that subterraneous fermentations, heat, and fire, worked many centuries to raise and accumulate them by many eruptions upon a calcareous marine ground, and perhaps in the midst of an ancient sea; but that water, rain, frost, and the inclemencies of the atmosphere since times immemorial, have been at work to destroy and to level them again.

Isolated, and for the most part conical ridges and points of basalt, of a like origin, and of the same remote antiquity, are dispersed

dispersed all over Hesse; and there seems to begin in Lower-Hesse in the neighbourhood of Cassell a chain of volcanic-hills, running through Upper-Hesse and the Wetterau to the Maya and to the Rhine, nay perhaps through Thuringia and Franconia to the Saxonian * and Bohemian mountains, which partly are known to be of a volcanic nature †. I would not however engage to prove, that, in the strictest sense, there are chains of volcanic mountains; however it is highly probable, that there is in Ger-

* For example, at Stolpen near Dresden.

† Baron Born's account of an extinguished volcano near Riga in Bohemia, printed in German, Prague, 1773; and Ferber's mineralogical description of Bohemia, Lusatia and Silesia are bordering to Bohemia, and Lusatia has a columnar basalt-hill near Grissenberg, as Silesia several others of that kind, which have been imperfectly described and drawn by Kundmann in rarioribus natura & artis. According to Baron Born's travels in Hungary, the Carpathian mountains are not destitute neither of that natural curiosity. I do not speak of similar observations made of late in Dalmatia, nor of those made in France.

many

many a long suite or succession of volcanos, comparable to that in Italy, which begins in the Southern skirts of the Alps in the Venetian State, runs on both sides of the Apennines into the kingdom of Naples, and ends beyond the Æolian and Liparean Islands in Sicily*. My conjecture is supported by these facts. Beginning from the Habichwald, there are on the N. W. side of the Fulde, through Lower- and Upper-Hesse up to Francfort, a great many isolated basalt-hills. Of that kind is the Densenberg in the bishoprick of Paderborn, nearer to the Habichwald and Dornberg in Lower-Hesse, the castle-hill at Grabenstine, the basalt-hills near Gudensberg, Felsberg, and Aldenburg, and those in the Wildenberg near Wolfhagen. In Upper-Hesse nearer to the Mayn there is a basalt-hill near Volckershausen, and another on the Vogelsberg at Bilstine †. They are conti-

* Ferber's Letters.

† Ritteri Tentamen historix naturalis Riedesellio-Avio-montanae in Tomo x. Actorum Societatis naturae curiosorum.

tion about Giessen. The castle of Amoneburg is situated on such a hill; Wilburg in Nassau likewise: Near Francofort on the Mayn, at Bockenhime, and near Andernac on the Rhine, the same volcanic tarras is found, which I have discovered on the Habichwald among many ancient traces of volcanic fire. But in the Northern parts of Germany; in Westphalia, the Electorate of Hanover, the Duchy of Brunſwick, the Hartz-Forest, the Bishoprick of Halberstadt and Magdeburg, the Duchy of Anhalt, and the Marches of Brandenburg, there have not been till this moment seen by me, or as far as I know found by others, any prismatic basalts or other unequivocal marks of volcanoes.

PART III.

UTILITY OF THESE ENQUIRIES, AND OF

THE VOLCANIC PRODUCTIONS.

General Remarks.

PEOPLE, to whom the very name and idea of a volcano is frightful, and to whom neighbouring volcanic ruins cause dreadful dreams of the earth-shaking contests of Pluto and Neptune, of the furnaces of Vulcan, or what is worse, of the eternal combustion of hell, objects and fancies by fools and poets seen in volcanos, will certainly bless these countries for being destitute of such troublesome objects. They will

will assure us perhaps, "that sleep and
 "life is surer where neither the marks of
 "ancient destructions are to be seen, nor
 "their returning rage to be apprehended.
 "Why will you force upon us old volca-
 "nos, unannoyed by history, and *good for*
 "*nothing?*" But there is much to be said
 to sooth such apprehensions. We live here
 on and near the ruins of our extinct vol-
 canes, as quietly and as securely as we
 should rest on the most bloody fields of an-
 cient battles, or on the tombs of raging ty-
 rants. Times immemorial their forces have
 been exhausted or quiet, and their present
 distance from the sea gives some hopes that
 they will be so for ever. They are besides
good for something, as shall be proved pre-
 sently; and this will justify the pains and
 care I have taken, and which other people
 may take, in their future examination and
 description.

I will not dwell much on the pleasure
 which enquiries of that kind afford to the
 mind

mind and the imagination; they are naturally bent to pry into the remotest antiquity, and into the first causes of things. But this pleasure, being personal, might perhaps be enjoyed, though the object of its enquiries be without any utility and advantage to human society. However, it may be a means of useful knowledge; and such is the conjecture and the conviction, that several of our mountains are of a volcanic origin.

§ 2. *Enquiries of this kind improve the knowledge of the physical geography, and the extensive art of the minerals.*

This knowledge improves and corrects our ideas concerning the origin and the natural vicissitudes of the surface of the earth; points out several dangerous errors, and teaches us to find at home several sorts of useful fossils, which either were not noticed at all, or were even at great expence imported from abroad—objects that certainly cannot

cannot be indifferent to a friend of truth, nor seem superfluous to a lover of his country, or to a man that once has felt the consequences of error! Every error is attended with its own natural punishment, and especially blunders committed in the expensive working of mines, whose punishment never fails to be immediate and extremely sensible. Such errors might be committed, and I am apprehensive have been committed very often, if in hope of metallic veins one should venture to sink shafts and to drive galleries through the vitreous rocks of volcanic mountains. They yield nothing but clear loss of money, pains, and time. Being accumulated by ashes, lavas, and straggling vomited stones, they may now and then contain in their melted masses and cinders some marks of metals; but their very nature forbids to hope or to look in them for metallic veins, which, by the subterraneous fermentation, heat, and fire, are destroyed and melted into one mass, with

the unweildy barren rocks that skirt them on every side. There has been under the late Landgrave Charles of Hesse, sunk a pit and a gallery through a basalt- and lava-rock, under the smaller Winterkaffen at the Habichwald. The gallery is still open, and is called the silver-well (silver-brunn). If these amazing subterraneous works have been undertaken on account of the cascades or the coal-pits, which are on the other side of the hill, they may perhaps be excusable; but if there has been any intention to fish for silver in the silver-well, as seems to appear by its name, the enormous expences have certainly been thrown away, and would have been saved by a better acquaintance with the nature of the whole mountain.

§ 3. *Use of the volcanic productions.*

I have given a description of the different earths and stones which have occurred to me in the Habichwald and other volcanic hills; and their number might perhaps
be

be augmented by such fossils as Mr. Ferber has found in the Italian volcanos; but, leaving this to future enquiries, I will at present dwell only upon the utility of the former.

§ 4. *Use of the basaltos and the harder lavas.*

The solid lavas and basaltos, on account of their remarkable hardness, are called *pietre dure** and *selce*†; and are employed in that country as well as in *Hesse*, 1. *in the paving of streets and roads*. Their plenty, hardness, and durable solidity, serves that purpose extremely well. They have been employed likewise, 2. *in walls of masonry*, as appears in the walls of many old castles and other pieces of old architecture in Germany; but, having been employed in their rough and irregular natural form, such walls are not to be commended. If the

* La Torre.

† Ferber, p. 230.

stones were cut into regular forms, it would make such walls extremely expensive, and in every other respect comparable to the everlasting Egyptian granite-architectures. In a natural prismatical form they serve at Felsberg, 3. as *posts on the street-corners* and, 4. as *palisades* to fence gardens and fields: They are set up close together, and thus make a fence or hedge of everlasting duration. However, they are seldom employed for this purpose, the carriage of the larger basalt-prisms being difficult and expensive. Some time ago the dispersed regular and prismatic basalt and lava pieces have been employed in the gardens at Wissemburg, 5. to border and to decorate some artificial brooks, ponds, natural cascades, and terraces; and to build occasional rustic chairs and stairs. Being very frequent within this noble garden on the descent of the Carlsberg, as well as in many picturesque and solitary valleys, they had been very troublesome to the gardeners, and rendered the clearing

clearing of the ground difficult and expensive, till the present gardener Mr. Swarzkopf, a man of taste and genius, had the happy idea to save time and expence, and to employ Nature's self to Nature's advantage. He composes these stones in whatever form he wants, or the situation asks and permits. Mosses and parasitical plants and shrubs serve instead of lime or cement. By this artifice he has produced a great many picturesque views, and many awful and singular objects, answering the wildness of their situation. Such are the rocky inclosures of Paul's-hermitage, in a lonely dale, surrounded on every side with hills and woods; the rivulets and cascades which from that valley, by many windings, and through many shrub- and tree-plantations, fall down to the Styx and the Chinese ponds; similar brooks under the Parnassus and Elysium on and near the Sneckenberg; and the steep rocky walls behind the Monuments of the Illustrious. You cannot
 imagine

§ 6. *Tarras and volcanic cinders employed by the Ancients in cast vaults and walls.*

Our tarras-stone can with great advantage serve in masonry, but especially in vaults. Being very light and spongy, its pressure and gravity is inconsiderable, and its coherence with lime and mortar inseparable. For this reason the Ancients and Italians have frequently employed this and the like sorts of stones. I shall prove what by the late *Abbé Winkelmann's* authority, and with his own words, which, in his *Remarks on the buildings and the architecture of the Ancients*, p. 9. runs thus:—"The Ancients, making their vaults very thick, wanted to have them as light as possible. This they effected by two means; the most common was, to build vaults with Vesuvian scoræ. These are reddish, or of a grey colour. A black species is dug out near Viterbo, in a place adjacent to some hot-wells. But these Viterbo flags being

" being not remarkably hard, are rather
 " unfit to be employed in vaults. The Ve-
 " suvian ones do manifestly appear in old
 " vaults. They have been occasionally ob-
 " served in the late reparation of the Pan-
 " theon. Vitruvius is silent about this
 " manner of vaulting; and he mentions
 " our Vesuvian slags but by the bye. The
 " nature of Vesuvius being scarce known to
 " the Ancients, they had neglected too the
 " examination of the Vesuvian products.
 " Vaults of such scoria are very common at
 " Naples; but Cardinal Alex. Albani has
 " made at Rome the first and till now only
 " experiment to revive this old practice, or
 " adding and exceeding such vaults in his
 " villa. The proceeding is thus; the cen-
 " tres of the scoria for the wall being
 " erected, they lay the arch on both sides
 " *lo scoria della volta* in the common way
 " up to the middle, and then pour or crown
 " it with scoria and mortar, which penetra-
 " ting into their spongy substance, makes
 " such vaults almost indestructible.

§ 7. *Example of a cast vaulting of calcareous
tuffas in Germany.*

Every light and spongy sort of stone would serve to the same purpose; and I remember to have seen at Lampring, an Abbey of English Benedictines in the Bishoprick of Hildesheim; very large vaults of a light calcareous tuffas or ostencolla. Necessity had laid them. The architect of the church had by ignorance or neglect treated its exterior walls so thin, that they were thought and found insufficient to support the weight of a stone vault. Therefore the church stood a long while unfinished; till at last they happily advised and executed a vault of this spongy calcareous substance; which without encumbering or destroying the exterior walls, supplies very well the want of a heavier one, though it should not be proof against bomb-shells, being everywhere penetrable even to the blade of a small sword.

§ 8. *Volcanic tarras gives an excellent cement.*

But the greatest, though in our countries most neglected, use of *tarras*, is its giving an excellent cement. I do not repeat what I have said about its volcanic origin and nature, which explains why it ought and might have the qualities of the Italian puzzolana.

§ 9. *The puzzolana described according to its volcanic origin, use amongst the Ancients, and its preparation.*

This puzzolana is an ash-powder, or rather pulverized cinders, found various-coloured about Naples, and black-coloured on *Veluvius* *. They dig it white and grey in the hills near *Puzzuolo*, brown and yellow at the foot of *Veluvius* †. It is likewise found near Rome in the before-mentioned

* Ferber, p. 147. † Ibid. l.c. p. 180, 181.

colour

colour and reddish*. It is produced by consumed cinders and scorified lavas; this is to say, by our tartas-stone; which resembles the scorified lava-fragments to be found in the Roman *puzzolana* †, for which reason our tartas-stone may be presumed to have the ferruminating and cementing quality of the *puzzolana*. This quality has been chemically analyzed by Mr. Ziegler ‡, and is, that *puzzolana* mixed with a convenient quantity of calcined lime-stone and water heats, and if heated and immediately employed as mortar, coagulates to such a degree of hardness and solidity, that water cannot dissolve or penetrate it §; nay, the mortar grows harder than even the stones which it is intended to bind. This appears in the bedding of the pavement of an ancient Roman-mosaico kept in the

* Ferber, p. 234, 235, 236. Winkelmann on the archit. of the Ancients, p. 6, 7, 8.

† See before.

‡ Magazine of Hanover, Nov. 21, 1774.

§ Ferber, p. 247.

grave's cabinet of antiquities. But the fairest proof to be given of it is the present state of the subaqueous ruins at Puzzuolo, Baja, and thereabout, and at Porto d'Anzio, where the pillars which inclosed the harbour are built of tiles. The Ancients employed the puzzolana likewise in the neighbourhood of Rome in the public roads; a practice not uncommon even in our days*. The black puzzolana is said to be excellent in buildings under water †, and the Ancients are said to have employed it on such occasions more than any other. ~~It is~~ other species duly mixed and prepared may answer the same purpose. For this reason this puzzolana is by large caravans exported from Italy into France, Holland, England, Sweden, and many other countries.

The preparation of puzzolana cement is as follows. Three or four parts of this

* Winkelmann's Remarks on Architecture, p. 7.

† Ibid. p. 6.

K.

substance

substance finely pulverized are to be mixed with one part of well-calcined limestone, likewise pulverized; then a good deal of fresh water is to be put upon that mixture, and the fluid mass to be turned and worked about till it begins to heat and to insipiate, when it is ready to be used and to be employed immediately; but care is to be taken that the walls and masonry made with puzzolana cement be covered for some time from sun-dry, and be kept wet by putting water upon them, else the cement will never coagulate entirely, and get in perfect solidity. It appears by this proceeding, that puzzolana cement cannot be properly applied to masonry, unless every piece of brick be sufficiently wetted, and the lines with which it is set, and that its sudden inspissation makes it naturally fit for incrustation, and expeditiously making large walls and vaults. The late Abbé Winkelmann describes it in the following passage, taken from his remarks on the architecture of the Ancients;

“ We

"We begin with the foundations, which
 "the Ancients laid either with large square
 "picce, or with small fragments of tufa.
 "The last method was the most common,
 "and is still used in Italy. They first dug
 "out the ground, and then filled the holes
 "by successively putting in puzzolana mortar
 "and tufa stones. Such a cement bed
 "coagulates in a couple of days, and gives
 "an excellent solid foundation. But it is
 "to be observed, that even in such masonry
 "above ground the Ancients employed
 "more cement than stone, which is a consequence
 "of its petrifying quality. Many
 "old vaults have been executed by the
 "same method. The scaffold having been
 "properly covered with wainscot of deal,
 "they put the above mixture of cement
 "and broken bricks and tiles upon it to
 "a determinate convenient thickness, which
 "in the Dioclesian Thermae is of nine palmi;
 "afterwards they covered and leveled the
 "whole by a bed of cement. By this
 "means large vaults could be finished in a

“ few days. That the Ancients commonly
 “ used it, is to be seen in old vaults; where
 “ they have given way, or lost their introp-
 “ tation, as in the Coliseo, in the Thermae
 “ of Titus, Caracalla, and Dioclesian, but
 “ especially in the vast ruins of the villa
 “ Hadriani, where even the impressions of
 “ the scaffold-deals are to be distinguished.”

This expeditious method of casting walls
 and vaults, is at present out of practice;
 however they continue in Italy to make
 their vaults with tufo and puzzolana, but
 as in common masonry; and then they
 strow and level the whole up to the back
 of the vaults with a castment bed, which
 casting they call *far a sacco*

PRO. The casting of walls and vaults by a
 puzzolana cemeny, used by the Ancients and
 the Italians, recommended.

This casting of foundations, walls, and
 vaults, supposes a plenty and low price of
 puzzolana,

puzzolana, and cannot be probably revived or introduced in countries where puzzolana must be imported; however, on account of its expeditiousness, which saves time and money, it certainly deserves great attention, wherever puzzolana is an inland commodity, or where, instead of puzzolana, there is to be found a tarras-stone, endowed with its ferruminating quality.

§ II. *The Hessian tarras proved by fact to give an excellent cement under water, and to be equal in goodness to the Dutch tarras.*

But here a question arises, “whether our Hessian tarras may be employed and worked as puzzolana? or whether it will give a cement in force and goodness equal to puzzolana cement?” This question has been answered to its advantage, not only by its origin and nature, but, what is above all reasonings *à priori*, by a great many experiments, which have been tried by me and

by Colonel Gohr, in several greater works under and above water at the Carlberg cascades at Weissenfline.

I hope, therefore, not only that our tarras may be henceforth conveniently employed in the country, but that it may be exported by the Weser, Carlshaven, and Bremen, to Holland and other Northern countries. For many years the tarras found near Bockenhime and Andernac on the Rhine has been exported in vast quantities to Holland, and thence to the North, though customs and duties make it very dear.*

The *ashes and friable cinereous stones* described above, may perhaps give likewise a good cement. They would sell somewhat

* The bushel of stamp tarras imported from Holland sells at London wholesale for three shillings or above. It might be had from Bremen at a cheaper rate, and in equal goodness, if ordered from the Hessian Company of Commerce established at Carlshaven on the Weser, who, applied to under this address, will supply any demands.

under

under the price of tarras, as not wanting to be pulverized, which causes some, though inconsiderable, expence. But, having made no experiments on their nature, I recommend them in the probable hope of their not being unsuccessful.

§ 12. *Mineral waters produced by the volcanic ashes in the garden at Weiffenfine.*

I mentioned before, that a cinereous stratum, under the Habichwald imparts to the water, which is soaking through it, a mineral and saline nature. Accordingly, it causes *mineral waters*, which on nearer observation I have found to be impregnated with some sal medium, vitriol, iron, and sulphureous acid. A well of such water appears under the bowling-green in the garden at Weiffenfine above the Styx, and another runs through the wall of a terrace before the gardener's house; both manifesting themselves by ochraceous sediments, and a volatile smell of hepar sulphuris.

But,


But, not being properly inclosed, and for that reason much adulterated by rain and other water, their mineral taste and mixture is weak at present, and will continue so till some famous and popular physician shall please to take this water under protection, and give it virtue and celebrity. Wells of a like, or even a different mineral nature, ought to be common round about the Habichwald, and other volcanic hills, especially in their cinerous strata.

The use of tufo being known; I conclude with wishing that these small discoveries may meet with a good reception, and be a recommendation of too much neglected Natural History.

THE END.

P. vi. l. 14. *for nomenclator read nomenclature.*

Plate I.
To face p. 16.



Handwritten text, mostly illegible due to fading and bleed-through from the reverse side of the page.

Basalte



Lefse.

*Plate II.
To face p. 38.*



Lava.



Books printed for L. DAVIS in Holborn,
Printer to the Royal Society.

1. **M**R. Da Costa's Natural History of Fossils,
upon a new System: To the Descrip^t
tions are added, Historical Facts and Critical
Observations, 4to. price 12s. 6d.

2. Mr. Meyer's Directions for making Mine-
ral Collections useful, 1s. 6d.

3. Fossilia Hartoniensia collecta, & in Museo
Britannico deposita, à G. Brander, *cum fig. ele-*
gantiss: & descript. à Solander, 10s. 6d.

4. Plukenetii Opera Bôtanica, Tab. æn: 434,
4 vol. 4to. 4l. 4s.

5. Dr. Halley's Tabulæ Astronomiæ, with
the Precepts in Latin & English, 4to. 1l. 1s.

6. Sir Isaac Newton of the Quadrature of
Curves and Analysis by Equation, with a Com-
mentary by Professor Stewart, 4to. 12s.

7. Newtoni Principia Mathematica, 4to. 12s.

8. Capt. Dobbs's Account of the Countries
adjoining to Hudson's Bay; their Situation, Cli-
mate, Trade, Navigation, &c. To which are
added, an Abstract of all the Voyages in the
Great Western Ocean, that have been published;
Vocabularies of Indian Languages; De Fontæ's
Voyage

Voyage from Lima; with other Pieces, also 35 in boards.

9. Peck's Memoirs of Milton, with critical Notes on Milton and Shakespeare, and several curious Pieces by Milton, &c. subjoined; R. P. 4to. 7s. 6d. sewed.

10. Memoirs of the Life and Actions of Greenwell, by John Milton; with a Collection of divers historical Desiderata, by Peck, and several engraved Heads, R. P. 4to. 7s. 6d. sewed.

11. Origines Typographicæ, à Meerman, 2 vols. 4to. 11. 4s.

12. Elstob's Saxon Grammar. With an Apology for the Study of Northern Antiquities, 4to. 2s. 6d. sewed.

13. Bibliotheca Literaria, a Collection of Inscriptions, Medals, and other Tracts, by Dr. Walfle and others, 4to. 6s.

14. Dr. Birch's Collection of Bills of Mortality, for a hundred Years back; with several curious Tracts subjoined on this Subject, 9s. In boards.

15. Travels through Part of Europe, Asia Minor, the Islands of the Archipelago, Syria, Palestine, Egypt, Mount Sinai, &c. Translated from the Low Dutch of the Hon. Mr. Van Egmond,

Egmont, Envoy to the Court of Naples, and Mr. Heyman, Professor of the Oriental Languages at Leyden, who have recorded many Particulars not mentioned by other Travellers, 2 vols. 8vo. 10s.

16. A Voyage to South America, from the Spanish of Geo. Juan, and Ant. De Ulloa, F.R.S. &c. &c. 2 vols. 8vo. 3d edit. 12s.

17. New Observations of two Swedish Gentlemen on Italy and its Inhabitants, from the French of M. Grofley, F. R. S. &c. &c. 2 vols. 8vo. 10s.

18. Mr. Grofley's New Observations on England, and its Inhabitants, made in a Tour to London in 1765; 2 vols. 10s.

19. The Beauties of England, or Traveller's Pocket Companion, 3s.

20. Dr. Goldsmith's Roman History, 2 vols. 8vo. 3d edit. 12s.

21. Dr. Goldsmith's Abridgement of his Roman History for the Use of Schools, 3s.

22. Baret's Travels in Spain, &c. 4 vols.

23. Father Charlevoix's History of Paraguay, 2 vols. 10s.

24. Falla's History of Jersey, 6s.

25. The

25. The History of China, upon the Plan of Rollin's Ancient History, 4s.

26. Floyd's Biography; or Lives of eminent Persons, British and Foreign, 3 vols. 18s.

27. Voltaire's History of Russia, with Maps, 2 vols. 10s.

28. An Etymological Celtic Vocabulary, 4s.

29. Parkin's Antiquities of King's-Lynn, and a Circuit of 15 Miles round, folio, 15s.

30. Badelade's Account of the Navigation of King's-Lynn, and of draining the Fens, with Charts, &c. folio, 12s.

31. Antiquities and Views of Egypt and Nubia, beautifully described, on 174 Copper-Plates, from Drawings taken upon the Spot, by Capt. Norden, R. P. fol. 11. 5s. half bound.

32. Capt. Norden's Travels in Egypt, with 174 Copper-Plates, 2 vols. 4l. 10s.

33. The History and Progress of Building, from the Creation to its Perfection in Asia, Egypt, Greece, and Italy, with 36 Plates, by J. Wood, of Bath, Esq; folio, 10s. in boards.

34. Dr. Warner's History of England, as it relates to Religion and the Church, 2 vols. fol.

35. Morant's Antiquities of Essex, 2 vols. fol.

